Benthic Community of Shuswap Lake's Foreshore

T.G. Brown, P. Winchell, and N. Postans

Fisheries and Oceans Canada
Science Branch, Pacific Region
Pacific Biological Station
Nanaimo, BC
V9T 6N7

2004

Canadian Manuscript Report of Fisheries and Aquatic Sciences 2693

Fisheries and Oceans Canada | Pêches et Océans Canada

Canada

Canadian Manuscript Report of Fisheries and Aquatic Sciences

Manuscript reports contain scientific and technical information that contributes to existing knowledge but which deals with national or regional problems. Distribution is restricted to institutions or individuals located in particular regions of Canada. However, no restriction is placed on subject matter, and the series reflects the broad interests and policies of the Department of Fisheries and Oceans, namely, fisheries and aquatic sciences.

Manuscript reports may be cited as full publications. The correct citation appears above the abstract of each report. Each report is abstracted in *Aquatic Sciences and Fisheries Abstracts* and indexed in the Department's annual index to scientific and technical publications.

Numbers 1-900 in this series were issued as Manuscript Reports (Biological Series) of the Biological Board of Canada, and subsequent to 1937 when the name of the Board was changed by Act of Parliament, as Manuscript Reports (Biological Series) of the Fisheries Research Board of Canada. Numbers 1426 - 1550 were issued as Department of Fisheries and the Environment, Fisheries and Marine Service Manuscript Reports. The current series name was changed with report number 1551.

Manuscript reports are produced regionally but are numbered nationally. Requests for individual reports will be filled by the issuing establishment listed on the front cover and title page. Out-of-stock reports will be supplied for a fee by commercial agents.

Rapport manuscrit canadien des sciences halieutiques et aquatiques

Les rapports manuscrits contiennent des renseignements scientifiques et techniques qui constituent une contribution aux connaissances actuelles, mais qui traitent des problèmes nationaux ou régionaux. La distribution en est limitée aux organismes et aux personnes de régions particulières du Canada. Il n'y a aucune restriction quant au sujet; de fait, la série reflète la vaste gamme des intérêts et des politiques du ministère des Pêches et des Océans, c'est-à-dire les sciences halieutiques et aquatiques.

Les rapports manuscrits peuvent être cités comme des publications complètes. Le titre exact paraît au-dessus du résumé de chaque rapport. Les rapports manuscrits sont résumés dans la revue *Résumés des sciences aquatiques et halieutiques*, et ils sont classés dans l'index annual des publications scientifiques et techniques du Ministère.

Les numéros 1 à 900 de cette série ont été publiés à titre de manuscrits (série biologique) de l'Office de biologie du Canada, et après le changement de la désignation de cet organisme par décret du Parlement, en 1937, ont été classés comme manuscrits (série biologique) de l'Office des recherches sur les pêcheries du Canada. Les numéros 901 à 1425 ont été publiés à titre de rapports manuscrits de l'Office des recherches sur les pêcheries du Canada. Les numéros 1426 à 1550 sont parus à titre de rapports manuscrits du Service des pêches et de la mer, ministère des Pêches et de l'Environnement. Le nom actuel de la série a été établi lors de la parution du numéro 1551.

Les rapports manuscrits sont produits a l'échelon régional, mais numérotés à l'échelon national. Les demandes de rapports seront satisfaites par l'établissement auteur dont le nom figure sur la couverture et la page du titre. Les rapports épuisés seront fournis contre rétribution par des agents commerciaux.

104-07560/

Canadian Manuscript Report of
Fisheries and Aquatic Sciences 2693

2004

BENTHIC COMMUNITY OF SHUSWAP LAKE'S FORESHORE

by

T. G. Brown[1], P. Winchell[1], and N. Postans[2]

[1] Fisheries and Oceans Canada
Science Branch, Pacific Region
Pacific Biological Station
Nanaimo, BC V9T 6N7
E-mail: browntg@pac.dfo-mpo.gc.ca

[2] Applied Technical Services
614 Bryden Court
Victoria, BC V9A 4Y5

© Her Majesty the Queen in Right of Canada, 2004
Cat. No. Fs 97-4/2693E ISSN 0706-6473

Correct citation for this publication:

Brown, T.G., Winchell, P., and Postans, N. 2004. Benthic community of Shuswap Lake's foreshore. Can. Manuscr. Rep. Fish. Aquat. Sci. 2693: iv + 41 p.

ABSTRACT

Brown, T.G., Winchell, P., and Postans, N. 2004. Benthic community of Shuswap Lake's foreshore. Can. Manuscr. Rep. Fish. Aquat. Sci. 2693: iv + 41 p.

Substrate trays and a benthic pump were employed to collect samples of the benthic community from the foreshore of Shuswap Lake, British Columbia. The majority of samples were collected in June and July (1999-2001) from depths of less than two meters. The most numerous benthic invertebrates were; Cladocera, Chironomidae (larvae), Ostracods, Oligochaeta, Calanoida and Nematoda. Although there was no significant diurnal difference in total invertebrate numbers, a greater number and weight of insects ($P < .05$) was measured from the samples taken at night in 2001. The cobble/rocky shores supported significantly more Cladocera ($P < .05$) especially chydorids and *Sida crystalline* than the sandy/mud locations. Slightly more chironomid larva were recovered from the sandy/mud foreshore locations, however this result was not significant.

RESUME

Brown, T.G., Winchell, P., and Postans, N. 2004. Benthic community of Shuswap Lake's foreshore. Can. Manuscr. Rep. Fish. Aquat. Sci. 2693: iv + 41 p.

Des échantillons de la communauté benthique de la zone intertidale du lac Shuswap en Colombie Britannique ont été recueillis en utilisant des plateaux de substrat et une pompe benthique. La plupart des échantillons ont été recueillis en juin et en juillet (1998-2001) à une profondeur de moins de 2 mètres Les invertébrés benthiques les plus nombreux étaient : Cladocères, Chironomes (larve), Ostracode, Oligochète, Calanoïdes et Nématodes. Malgré que l'effet des variations diurnes sur le nombre total des invertébrés n'était pas significatif, un plus grand nombre d'insectes ayant un poids plus élevé ($P > .05$) a été mesuré à partir des échantillons prélevés la nuit en 2001. Les rives rocheuses/rocailleuses accueillent un plus grand nombre de cladocères ($P < .05$) surtout les espèces chydoridae et Sida crystallina que les endroits sableux/boueux. Un nombre légèrement supérieur de larves de chironomes a été recueilli dans la zone intertidale aux endroits sableux/boueux, cependant ce résultat n'est pas significatif.

TABLE OF CONTENTS

ABSTRACT III
RESUME III
INTRODUCTION 1
STUDY AREA 1
METHODS 1
RESULTS 3
DIFFERENCES IN METHODS AND YEARS 3
INVERTEBRATE NUMBERS AND SEDIMENT 4
INVERTEBRATE NUMBERS VS SUBSTRATE AND RIPARIAN TYPES 4
NON-INSECTA GROUPS 4
Ostracoda 4
Acari 5
Copepoda 5
Cladocera 5
Nematoda 6
Mollusca 6
Amphipoda 7
Oligochaeta 7
INSECT NUMBER VS WEIGHT (2001) 7
NIGHT VS DAY FOR INSECTS 7
INSECTA GROUPS 8
Diptera 8
Coleoptera 9
Collembola 10
Trichoptera 10
Heteroptera 10
Hemiptera (Aphididae) 10
Ephemeroptera 11
DISCUSSION 11
ACKNOWLEDGMENTS 13
INTERNET SOURCES 13
REFERENCES 13
LIST OF FIGURES 15
FIGURES 17
APPENDIX 1. BENTHIC SPECIES LIST; SHUSWAP LAKE FORESHORE 29
APPENDIX 2. SHUSWAP LAKE BENTHIC SAMPLES (1999-2001) 34

INTRODUCTION

Shuswap Lake is an important source of Thompson River sockeye (*Oncorhynchus nerka*), chinook (*O. tshawytscha*), and coho (*O. kisutch*) salmon. Juvenile salmon utilized the foreshore areas of Shuswap Lake for rearing and migration (Fedorenko and Pearce 1982; Graham and Russell 1979; Russell et al. 1980). Juvenile salmonids can exploit previously dry lake margins (Figure 1) from April to July, as the lake water levels rise. Fish can also access alcoves, channels, and bordering wetlands previously isolated from the lake. A large portion of the diet of juvenile coho and chinook salmon captured from the foreshore of Shuswap Lake consisted of benthic invertebrates (Brown and Winchell 2002).

The foreshore (littoral zone) of large lakes such as Shuswap Lake is under considerable pressure from urbanization. Some beaches are naturally sandy (Figure 1), but cobble and boulder substrates cover much of the foreshore (Figure 2). Cobble substrate is removed for private beach creation and collected in rows, riparian vegetation is cut, protective walls are built, emergent vegetation is removed, and small marshy bays are altered. We studied the littoral zone associated with the main arm of Shuswap Lake for 3 years (Figure 3). Field-work was conducted during spring and summer (April to August) as lake water levels rose and fell. This study was designed to define the role of Shuswap Lake's littoral zone as juvenile salmon fish habitat and assess the potential impacts of foreshore and riparian development.

This paper examines the benthic community of Shuswap Lake's seasonally flooded foreshore. An examination of the benthic community is only a small component of the overall ecology of juvenile salmonids rearing along the foreshore of a large lake. Fish diets and distribution will be examined in a later paper.

STUDY AREA

Shuswap Lake is located in the southern interior of British Columbia within the South Thompson River drainage. The South Thompson River drains an area of approximately 16,200 km^2 (Fedorenko and Pearce 1982). Shuswap Lake is a multibasin lake consisting of four arms or five reaches (Salmon, Sicamous, Anstey, Seymour, and Main), has a surface area of 310 km^2, has a maximum depth of 162 m, and is considered to be oligotrophic (Williams et al. 1989). Lake transparency as measured mid-lake near Sorrento can range from 2.0 m during the period of peak runoff to 12 m in winter or early spring (Williams et al. 1989). The Main Arm of Shuswap Lake was the only area examined during the three years of study (Figure 3).

Shuswap Lake water levels were monitored at Sicamous B.C. by Kramer (2003) and were recorded as meters above sea level. The lowest Shuswap Lake water levels occur in March (e.g. March 22, 2001 at 344.8 m elevation) and the highest lake levels have historically occurred in June or July (e.g. June 14, 1972 at 349.66 m elevation). Lake levels annually rise from 3 to 4.5 m freshly flooding a considerable portion of the lake margins and occasionally inundating surrounding riparian vegetation (e.g. 1997 and 1999). The largest single day increase in lake level was 24 cm on May 28, 1972. During the three years of study the largest single day increases in lake levels occurred on; May 26, 1999 at 16.0 cm, May 22, 2000 at 12.6 cm, and May 28, 2001 at 16.1 cm. The maximum recorded lake level for each of the three years of study were; 349.22 m on July 15, 1999, 348.33 m on July 1, 2000, and 347.92 m on June 11, 2001 (Kramer 2003). Davies et al. (1996) reported an average annual difference in water levels of 3.3 m, with a mean high water of 348.3 m above the geodetic chart datum and a mean low water level of 345.0 m.

METHODS

Benthic samples were collected from the littoral zone of Shuswap Lake for three years. Different methods and study design were employed each year and it is likely that each sampling method was biased towards the collection of different invertebrate groups. The difficulty of sampling cobble bottoms

with the same efficiency as sand/mud bottoms excluded the use of "dredges and grabs" (Southwood 1966) and led to the planting of removable portions of the substrate within plastic trays in 1999 and 2000. Attempts to measure the types and production levels associated with different substrates, using plastic trays were unsuccessful during the first two years. In 2000, the majority of substrate baskets and the invertebrates that had colonized them were never recovered. A benthic sampler was used to vacuum invertebrates from the shallow lake foreshore during the last year of study. This benthic pump could only sample shallow depths (< 0.4 m). Thus, as different methods were used each year, it is difficult to compare years.

In 1999, three plastic benthic trays were placed at each of 16 locations (Appendix 1, Table 1-2). Eight locations were characterized as having sand/mud beaches and eight had cobble beaches. Each tray (23 x 15 x 5 cm) covered an area of 345 cm^2 and contained approximately 1725 cm^3 of substrate collected from the immediate foreshore (Figure 4). All larger cobbles and rocks were removed (> 8 cm) and the trays were buried 15 cm into the foreshore (level to the top of the tray). Trays were installed on June 8-10 at a depth of 50-70 cm and at 75-90 cm on June 17-20 as lake levels rose. After lake levels had peaked and fallen the trays were recovered on July 28-30 from a depth of 50-95 cm. Three trays from Location 8 could not be found and 3 grab samples of a similar size were substituted for the missing tray samples at the time of recovery.

In 2000, two locations were chosen for benthic collection (Appendix 1, Table 3). Thirty of the same benthic trays used in 1999 were used in 2000; however, all trays contained a uniform 1725 cm^2 of sieved gravel 3.2 – 6.3 mm (1/8 to 1/4 inch). Both locations had been altered by the clearing of boulder/cobble material from portions of the foreshore and placement of these rocks into rows (Figure 5). Each location was divided into 3 substrate types; natural cobble/rock, sand/fine-gravel following cobble removal, and windrowed cobble. At each location 5 trays were placed within each of the three substrate types. This study design allowed the comparison of windrowed cobble and natural foreshore. The trays were placed on June 4 at a depth of 60-80 cm and after lake levels peaked the trays were recovered on July 31 from 40-60 cm depth. Unfortunately only 10 of the 30 trays were recovered.

The trays and contents were separated from the surrounding bottom materials, placed in a bucket under water, and hauled to the water surface. All fine materials and invertebrates held within the bucket were then washed and sieved in a fine mesh net (200 micron nitex) to reduce water volume. Concentrated materials were emptied into a 1 liter jar and preserved with formaldehyde. To estimate the number and type of invertebrates that could potentially be added to the sample by this removal method, a series of six bucket hauls with an empty tray were completed.

In 2001, a total of 37 samples were taken by a customized benthic pump sampler (Appendix 1, Tables 4-6). The first test sample was taken on April 26. To compare night and day benthic activity a series of 3 night and 3 day samples (30 samples total) were taken at 5 locations from May 10 to August 28. Six additional daylight samples were taken from two locations during peak flooding (June 19 to July 11).

The benthic pump sampler vacuumed benthic organisms and fine materials from the shallow edge of Shuswap Lake (Figure 6). This apparatus consisted of a bilge pump and associated hoses, a portable power supply, a modified circular "Hess" type benthic frame sampler with fine-mesh screening (300 micron), and fine mesh collecting net (200 micron) with screw sieve collecting container (200 micron). The frame of the sampler was 40 cm deep and 43 cm in diameter. The area sampled was 1450 cm^2 and during sampling the bottom was turned over to a depth of 3-4 cm. This equates to a substrate volume of approximately 5000 cm^3. Water depth ranged from 10 to 27 cm and mean water depth sampled was 17.8 cm (± 1.2 cm, 95% Confidence Limits or C.L.).

The benthic pump sampler was designed to sample both cobble and fine substrates in a similar manner. However, when the sampling frame is being placed on cobble beaches, larger stones will lie partly inside and partly outside the metal frame and selection and rejection of these materials is a matter of judgment. The larger rocks considered inside the frame were rubbed with gloves (inside of frame) and removed from the sampling area. Following removal of all larger rocks, the top 3-4 cm of the remaining

finer substrate was stirred by hand and the materials and water within the frame was vacuumed for 10 min. The collecting net (200 micron) was washed (filtered water) and benthic invertebrates and fines were concentrated in the removable plastic sieve end of the net (200 micron). The concentrated materials were emptied into a 1 liter jar and preserved with formaldehyde. All invertebrates were identified and counted and all insects in 2001 were weighed (Appendix 1, 2).

Location 7 was sampled both day and night in early spring 2001 as the lake level rose and again in August 2001 as lake level fell. The riparian zone at this location consisted of a grove of cottonwoods and willows and the substrate consisted primarily of sand and muck. The foreshore had a gentle slope and in June/July the lake flooded into this riparian vegetation as well as covered small patches of emergent vegetation (mainly grasses and sedges). We considered this location to be undisturbed by lake front development.

RESULTS

DIFFERENCES IN METHODS AND YEARS

The method of benthic sampling was different during each of the three years. In 1999, substrate trays were filled at each location with unwashed foreshore substrate materials representative of that location. Samples obtained in this manner would approximate samples obtained with a benthic grab. Invertebrates, invertebrate eggs, fines, and detritus present at the location would be added to the tray at the time of filling. That portion of the invertebrate community that utilized fines and detritus and is slow to colonize new areas, should be present in greater relative numbers (Figure 7).

In 2000 all fines, organic materials, and invertebrates were washed from the sorted gravel used to fill the trays. Every tray held the same gravel mix. This was considered to be an "artificial" substrate as it was not representative of the foreshore substrates upon which it was placed. Those invertebrates that could rapidly colonize the trays and that preferred cleaner substrates should dominate (Figure 8).

In 2001 a benthic pump was used to sample the shallow (<40 cm) freshly flooded portion of the foreshore. This method should be biased towards two types of invertebrates; those that rapidly colonize the flooded foreshore in May and June and those that remain in the shallow waters as the lake levels drop in July and August (Figure 9). In 2001, sampling commenced in early May instead of June-July as in 1999-2000 and 15 night samples were taken. Thus, those invertebrates more prevalent on the shallow lake margins at night and in early spring would be relatively more numerous in the 2001 samples.

During the three years of study a total of 114,574 invertebrates were classified (Appendix 1; Figure 10). The most numerically abundant benthic invertebrate group was Cladocera 27,212 and it was also the least variable, annually averaging 23.8% of the invertebrates identified (Figure 11). The other dominant invertebrate groups included; Chironomidae (19.1%), Ostracoda (17.5%), Calanoida (11.0%), Oligochaeta (13.6%), and Nematoda (8.7%). These six groups represented 94% of the total benthic numbers (Figures 10-11).

Differences in the relative composition of the benthic community were noted for each year of the study (Figure 11). In 1999, Oligochaeta and Nematoda were 10x more common than in 2000. These worms utilize fines and detritial material and we suspect they were placed in the trays with the unwashed substrate, but were slow to colonize the cleaned substrate used in 2000. Calanoida and Chironomidae larvae were twice as prevalent in the community structure in 1999 as in 2000. These invertebrates also rely on detritus and may have been reluctant to colonize the cleaned substrates. Ephemeroptera (mayflies) and Trichoptera (caddis flies) dominated the clean gravel trays used in 2000 compared to the unwashed substrate used in 1999.

The benthic pump samples did appear biased towards Ostracoda, capturing 30 times more in 2001 than the benthic trays in 1999 or 2000. We observed large numbers of very small highly mobile invertebrates in very shallow water at night in April-May. Thus the higher numbers noted in 2001 are

likely due to when the samples were taken and not necessarily due to the efficiency of the pump sampler compared to tray samples. Nematodes were also relatively more abundant in 2001 samples. Although nematode food and feeding habits vary among freshwater groups, many of the nematodes we captured could be considered to be microbiotrophic (feeding on bacteria, algae, fungi, and protozoa). Decaying vegetative material, concentrated along the shallow lake edge (wrack), may have provided a good food source for both Ostracoda and Nematoda.

INVERTEBRATE NUMBERS AND SEDIMENT

The volume of fine material remaining (< 200 microns) after the samples were sieved was measured for each sample in 1999. This volume of fines can be compared to the total number of invertebrates (Figure 12) and to the number of non-insecta invertebrates (Figure 13). There was a positive correlation) between the volume of fines and the total number of invertebrates (Petersons $P < .05$). There was also a positive correlation between the volume of fines and the number of non-insecta invertebrates (Petersons $P < .05$). Although the number of each non-insect invertebrate group was not significantly correlated to volume of fines, the actual number of ostracods, oligochaetes and nematodes increased with the volume of sediment.

INVERTEBRATE NUMBERS VS SUBSTRATE AND RIPARIAN TYPES

In 1999, three trays were placed on each of 8 cobble/rock foreshore locations and 8 sand/mud locations. A difference between the total number of invertebrates and substrate type was not apparent. The total number of benthic organisms was similar under the two treatments (Figure 14). In 2000, not enough trays were recovered to reasonably analyze the data. However, it appears the number of organisms was similar on both of the substrate types (Figure 15).

In 1999, the riparian zones of 8 locations were considered to be natural or treed (supported well established natural trees and brush) and 8 locations were considered to be cleared (majority of riparian zone consisted of lawns, riprap, retaining walls, roadways, and developed lands). There was no apparent difference in total invertebrate numbers when the two riparian types are compared. Possible differences in the abundance of invertebrate families relative to habitat types will be examined below.

NON-INSECTA GROUPS

Ostracoda

The relative number of each invertebrate group changed over the season. In temperate lakes, many of the Ostracoda occupy the shallow edge of the lake (<30 cm depth) as the water level rises in May (Figure 16). These benthic invertebrates reside on the sediment-water interface of the substratum feeding on small pieces of detritus and algae (Delorme 1991). They are very small (< 1.5 mm) and the majority of the body is contained within a solid carapace (Walter 2002). These Ostracoda species are not tolerant of warm summer temperatures (Delorme 1991) and few were collected in July and August from the warm foreshore.

Three Ostracoda genera dominated the foreshore of Shuswap Lake. *Cypria* sp and *Cypris* sp. were significantly more numerous in May than in August (Two-Way ANOVA, $P < .05$) and significantly more numerous at night (Two Way ANOVA, $P < .05$). The spring numbers of these two genera represented 92% of the total number of Ostracoda captured from all locations during 2001 (Appendix 1, Table 4). In 1999, *Candona* was the more numerous genera captured (Appendix 1, Table 1); while in 2001 they were the third most common genera. In 2001, *Candona* sp. were more numerous during the day but not significantly.

Acari

Among the Acari, the Hydracarina are commonly referred to as "water mites" (Ward and Whipple 1966) and most have three life stages; larvae, deutonymph, and adult (Clifford 1991). The larval stage is parasitic, commonly as an ectoparisite on immature aquatic insects (Smith and Cook 1991). Only free-living adults were enumerated during this study. Hydracarina were more numerous on rocky shores than on sandy/mud shores in 1999 (t-test P <.01), 2000 (t-test P <.05), and from the samples collected in June and July 2001 (t-test, P < .01). The family Oribatidae includes the genus, *Hydrozetes*. These adult mites are herbaceous, feeding on fungi and green plant material (Ward and Whipple 1966). Location 7 supported beds of aquatic vegetation. This location (Figure 16) also supported the most Oribatidae (100/283 in 1999 and 230/284 in 2001).

Copepoda

Copepods are common in a wide variety of aquatic habitats (benthic, littoral and pelagic) and most species are omnivorous (Williamson 1991). They may feed on detritus, pollen, plankton, other invertebrates and even larval fish. Williamson (1991) described the calanoids as primarily planktonic and the cyclopoids as primarily littoral and benthic. Adams et al. (1990) noted that the most dominant epibenthic species in Shuswap Lake were Copepoda (Calanoida and Cyclopidae).

Although Copepoda were found at all locations, the majority (1055/1729) in 2001 were found at sites supporting aquatic vegetation (location 7; Figure 16). In 2001, they were more abundant later in the year (80/sample) compared to early in the year (25/sample) but not significantly so (t-test). The most numerous Copepoda suborder was Cyclopoida (6642/10755 in 1999, 91/169 in 2000, and 1565/1729 in 2001).

Cladocera

The Cladocera (water fleas) are a group of small (range from 0.2 to 18 mm) meiobenthic crustaceans (Dodson and Frey 1991) and are a common item in Shuswap Lake fish diets (Russell et al. 1980). The Cladocera consume a variety of small particles ranging from bacteria, through algae to rotifers and copepod nauplii. We identified six families of Cladocera of which the Chydoridae and Sididae were by far the most numerous (Appendix 1, Tables 1,3,4). The Cladocera were more abundant on rocky foreshore than on sand sites in 1999 (Mann-Witney P < .05). Cobble/rocky locations averaged 432 ± 95 (95% C.L.) cladocerns / sample while sandy/mud locations averaged 194 ± 45 (95% C.L.) cladocerns / sample.

The Daphnidae (*Scapholeberis rammneri* and *S. mucronata*) are primarily limnetic (Edmondson 1966). The species in the genus *Daphnia* are limnetic zooplankton but may be encountered in the littoral zone (Chengalath 1982). *Ceriodaphnia* are small in size but were common in littoral samples collected across Canada (Chengalath 1982). They range in size from 0.2 to 3.0 mm and most are filter feeders using setae on 5 pairs of legs to capture small particles (Holmes 2001). Very few Daphnidae were captured in the substrate trays used in 1999 and 2000. In 2000, more Daphnidae were found in the six water column samples than in the 10 substrate samples. This indicates they reside in the water column not on the bottom. In 2001, three of seven samples taken in April and May had 95% (770/807) of the total number of Daphnidae. This indicates a highly gregarious distribution and possible higher spring abundance of Daphnidae on the lake foreshore. Their occurrence on the lake foreshore is likely related to winds and lake currents rather than habitat preference.

The Chydoridae (family of Cladocera) are typically littoral (Chengalath 1982). In 1999, three chydorids dominated the foreshore in July. These included; *Acroperus harpae*, *Alona* spp., and *Chydorus sphaericus* (Appendix 1, Table1). *A. harpae* were significantly more numerous on rocky foreshore than on sandy/mud foreshore in 1999 (Mann-Whitney P < .0001). Chengalath (1982) reported that *A. harpae* was common to all lake habitats. Edmondson (1966) reported that it prefer shallow, weedy habitats and was rare in open waters. *Pleuroxus procurvatus* and *P. denticulatus* were found at only a few locations but these were sand/mud bottomed. *Alonella* spp. were more numerous at two cobble/rock locations

(610/721) and *Camptocercus rectorostris* was more abundant at cobble/rock locations (496/640). In 2000, the chydorids were more numerous within samples obtained from cobble/rock sites than from sand/mud sites (t-test, $P < .05$).

The chydorids were significantly more abundant along the shallow foreshore as lake levels dropped in August 2001 than in May (t-test, $P < .01$; Figure 16). *Alonella excisa* and *Chydorus sphaericus* were 100 times more numerous in August than in May. *Acroperus harpae* was more prevalent in June and slightly more were captured during the day than at night (Appendix 1, Table 4).

Polyphemidae (*Polyphemus pediculus*) is common in marshes and weedy ponds (Edmondson 1966). In 1999, it was found only at location 7. This location contains considerable emergent vegetation (reeds and sedges).

Sida crystallina were more numerous on rocky-cobble foreshore in July 1999. (Mann-Whitney $P < .0001$). In 2000, *Sida crystallina* dominated the Cladocera (541/604) and were more numerous at the cobble/rock locations than at the sand/mud locations (Mann-Whitney, $P < .05$). A few *Sida crystallina* were captured during the control sampling in 2000. This indicates they were common in the water column as well as on the lakeshore bottom. These organisms should be readily available to juvenile fish feeding in the littoral zone. In 2001 during June and July, significantly more *Sida crystallina* were captured in benthos samples obtained from cobble/rocky shores than from sandy/mud sites (Mann-Whitney, $P < .001$).

Nematoda

Nematoda are more likely to be captured through the use of a Galen type sampler (samples substrate) than by a benthic sled (Adams et al. 1990). They tend to occupy the interstitial spaces in the substratum rather than the surface. It is doubtful that they are available as juvenile salmonid food. The Nematoda (Figure 17) were more abundant in May than in August 2001 (t-test $P < .01$). Nematoda were also more numerous in the 1999 benthic samples with a greater volume of fines; although the correlation between sediment and numbers was positive, it was not significant (Peterson, $P = .09$).

Mollusca

The bivalves (Pelecypoda) are filter feeders, consuming plankton, detritus and bacteria. They are called "fingernail clams" (Soil and Water Conservation Society of Metro Halifax 2002). The freshwater Sphaeriidae are often abundant in profundal areas, residing in the sandy substrate. Two genera were noted (*Sphaerium* and *Pisidium*). *Sphaerim* sp range in size from 8-20 mm, while *Pisidium* sp are smaller, ranging in size from 2-6 mm. It appears the bivalves were highly gregarious in distribution. In 2001, all 209 of the bivalves identified were obtained from location 7. No bivalves were recovered from the plastic substrate trays in 1999 or 2000.

The freshwater Gastropoda (snails) are primarily herbivores and detritovores. They use a rasping radula to scrape periphyton from substrate and epiphytic algae from macrophytes (e.g. *Gyraulus*), or consume macrophytes (e.g. Lymnaeidae). The genera we captured have been described as small, lentic, thin shelled, and preferring sandy to muddy substrates (Schuster 2002).

Gyraulus spp. were present each year and were the dominant gastropod (134 of the 166 characterized to genus). *Stagnicola* and *Physa* spp. were also found, but in fewer numbers. In 1999, significantly more gastropods were found on the cobble-rocky shores than the sand/mud beaches (t-test $P < .05$). In 2000, *Gyraulus* sp. was more abundant in the 2 samples taken from sandy substrate (36/sample) than in the 5 samples taken from cobble substrate (7.2/sample) at location 14. However, as few samples were recovered, statistical analysis is impossible. In 2001, more gastropods were collected on the sandy shores (1.2/sample) than on rocky locations (0.67/sample) although this result was not significant (t-test, $P = 0.28$).

Amphipoda

The Amphipod, *Hyalella azteca* is a relatively large laterally compressed invertebrate (> 6 mm), easily seen in the shallow lake waters and is commonly referred to as a "scud." It is an omnivorous scavenger, feeding mainly on dead animals and plant matter (Clifford 1991). *Hyalella* requires well oxygenated water, resides in water less than 1 meter deep, and often hides under plant mats. In 1999, 200 of 223 *Hyalella* were captured from rocky foreshore sites. In 2000, 8 of 10 were from the trays placed on cobble foreshore. In 2001, more than half (589/983) were recovered from location 7 and an additional 230 were enumerated from the single sample taken at location 2 in April. At the time of sampling location 2, wave action had covered the foreshore with a blanket of fine decaying organic materials (leaf litter and wood chips).

Oligochaeta

Another common group of invertebrates identified within the benthic samples from Shuswap Lake was Oligochaeta (Figures 11, 17, 18). Three families dominated, these were Naididae, Enchytraeidae, and Tubificidae (Appendix 1. Tables 1,3,4). The freshwater oligochaetes are primarily burrowing worms, feeding by passing mud and debris through the gut and extracting organic matter (Clifford 1991). It is doubtful that oligochaetes are available to juvenile salmon. These worms are generally found in soft sediments rich in organic matter and they can tolerate low oxygen levels (Fox 2000). In Shuswap Lake, some naidids and tubificids are epiphetic (Hatfield 1996). *Nais* sp. feeds on heterotrophic aerobic bacteria, while *Chaetogaster* spp. are carnivorous, actively hunting and consuming smaller organisms such as ostracods (Nikon Microscopy 2003). The enchytraeids are common in marginal aquatic habitats such as marshes, small streams and interstitial waters along the margins of lakes (Soil and Water Conservation Society of Metro Halifax 2002).

In both 1999 and 2001, the naidids dominated the counts of identified aquatic oligochaetes, representing 91% and 74% of the oligochaetes. In 2000, only 13 worms were obtained from the cleaned substrate trays (Appendix 1, Table 3). The most commonly identified species of naidids were; *Chaetogaster* spp., *Nais* spp. *Stylaria lacustris* and *Pristina foreli* (Appendix 1, Table 1, 4).

The distribution of oligochaetes was location specific in 2001 (Figure 18). The sites with the most flooded aquatic vegetation (locations 2 and 7) had the highest densities of oligochaetes, mainly naidids. However, high densities of tubificids were noted in August at location 7. Location 10 also had a high density of enchytraeids. There was not a significant correlation between the volume of sediment/sample and the number of oligochaetes/sample in 1999 (Linear Regression $P = .15$). In 1999, there was no significant difference between the number of oligochaetes within cobble/rock and sandy foreshores (t-test, $P = .72$).

INSECT NUMBER VS WEIGHT (2001)

An assessment of the number of insects without consideration of their relative size is misleading. The midges (Chironomidae) represented 59.1% of the total number of insects in the 2001 benthic pump samples (Figure 19). The combined numbers of caddisflies (Trichoptera), beetles (Coleoptera) and true bugs (Hemiptera) represented only 9.4% of the total number of insects. When the weight of the individual insects is examined (Figure 20), the latter group represented 50% of the total insect weight and the midges only 20.8%. In terms of numbers, the midges and mayflies (Ephemeroptera) dominate. However, four groups (caddisflies, mayflies, midges, and true bugs) were similar in relative group weight and cumulatively represented 86% of the total insect weight.

NIGHT VS DAY FOR INSECTS

There were significantly more (2 Way ANOVA $P < .05$) insects at night (139.8/sample ± 18.0 1SE) versus day (94.0/sample ± 13.1 1SE) in 2001 (Appendix 1, Table 5). The insects were more numerous at two locations in 2001 (2 Way ANOVA $P < .001$). Locations 10 and 7 supported higher numbers of insects than the other sites, mainly due to the high numbers of Chironomidae recovered at

those two locations. Location 7 was sampled both day and night (Figure 21), while location 10 was not sampled at night.

The weight of insects / sample was greater at night (87.4 mg/sample ± 33.18 1SE) than during the day (26.4 mg/sample ± 8.65 1SE), but not significantly so (Mann-Whitney P = 0.15). Location 7 represented 94% of insect weight from comparable night/day samples in 2001 (Figure 22). If we compare the day/night insect weights from only location 7, then a significantly greater weight of insects was obtained from the night samples (Mann-Whitney P < .01). This apparent weight difference is due to a few large Trichoptera, Corixidae, and Coleoptera captured at night (Appendix 1, Table 6).

INSECTA GROUPS

The midges (Chironomids) were the most numerous insect group identified during this study (Figure 11; Appendix 1, Tables 2, 3, 5). They represented 87% of the total insect number and annually represented; 96.7% in 1999, 40.4% in 2000, and 59.1% in 2001. The next most numeric group of insects were the mayflies (Ephemeroptera). They represented 8.1% of the total insects (1999, 1.6%; 2000, 51.3%; 2001, 25.5%). Together the midges and mayflies represented 95% of the insects recovered from the foreshore of Shuswap Lake during the three years of study. The relative numbers of the remaining Insecta groups changed slightly during each of the three years (Figure 11) with Diptera other than Chironomids (1.8%), Trichoptera (1.6%), Coleoptera (0.5%) and Collembola (0.3%) the next most abundant groups.

In examination of the foreshore invertebrates, two groups of Insecta common to freshwater were not found. These are Culicidae (mosquitoes) and Plecoptera (stoneflies). This is not surprising as mosquito larva must obtain oxygen at the lake surface and large lakes have considerable wave action making this difficult or impossible. Stonefly nymphs require well oxygenated water. Consequently they are found in flowing rivers and streams amongst the rocks and bottom debris. A few species of stoneflies can also be found in the rocky shoals of cold lakes.

Diptera

Four families (Chironomidae, Ceratopogonidae, Dolichopodidae, and Empididae) were numerically the most dominate of the "true flies." The Chironomids were the most numerous and were found at all sites. All non-chironomidae were captured as larvae and appeared to be very location specific rather than spread uniformly around the lakeshore.

Chironomidae

The Chironomidae are one of the most important food sources for salmonids (Newman 2003). The pupae are the most vulnerable to fish predation as they must secure oxygen from the water surface through an anterior respiratory horn. The larvae dominate the benthos within the littoral zones of a lake and their faecal pellets are an important component of the food chain. The larvae are primarily herbivores or detritovores, consuming fine particles and blue green algae. Many of the larvae construct sediment tubes in the substrate and are filter feeders. The pupae and adults do not feed. The adults fly over the water and may temporarily land on the surface.

In 1999, we estimated the density of chironomid larvae to be greater than 1/cm^2 (Figure 23). In later years the density was substantially less. It is possible this difference was due to annual changes in sampling procedures as described earlier. However, it could also be due to differences in sampling locations or seasonal flooding.

Considerable variation exists between sampling locations for any given year (Appendix 1, Tables 2, 5). There is too much variation between locations to clearly show differences in larval densities relative to the physical nature of the foreshore. Slightly more chironomid larva were recovered from the sandy foreshore locations (414/sample ± 121, 1SE) than from the cobble beaches (379/sample ± 121

1SE) in 1999. However, one sandy location (location 8, grab samples) strongly biases this result. If location 8 is excluded from the analysis then the cobble locations have significantly (Two Way ANOVA, $P < .05$) more larvae than the remaining sandy locations (150/sample ± 60 1SE). The treed foreshore locations (281/sample ± 60 1SE) had a similar number of chironomid larvae to the cleared locations (249/sample ± 63 1SE) in 1999. In 2000, more chironomids were recovered from the trays placed on cobble substrate (42/sample ± 10 1SE) than from trays placed on sandy substrate (18/sample ± 6 1SE). This result was not significant (t-test, $P = .11$). In 2001, cobble locations supported 15 larvae / sample (± 7 1SE) versus 102 larvae / sample (± 22 1SE) on sandy foreshore locations. This result is also strongly biased by the high density noted for sandy location 7 (195 ± 22 1SE or 0.13 / cm^2).

More chironomid larvae were captured at night than during the day (Figure 24). However, there was not a significant difference between day and night larval densities (Two-Way ANOVA $P = .15$), while there was a significant difference between larval densities at location #7 compared to the other three locations ($P < .001$).

A portion of the Chironomidae captured during the three years of study (15,793 of 21,894) was classified further. Three subfamilies dominated the Chironomidae found along Shuswap Lake's foreshore. The most numerous were Chironominae of which 6,980 were of the tribe Tanytarsini and 1,828 were of the tribe Chironomini. Orthocladiinae numbered 5,448, Tanypodinae numbered 518 and 18 Diamesinae were identified.

Ceratopogonidae

Some adult ceratopogonidae bite warm blooded animals and these are commonly called "biting midges" or "no-see-ums." Others such as Bezzia and Probezzia feed on small insects (i.e. Chironomidae). As larvae they are small, elongate, carnivores, and swim with a serpentine motion (Hilsenhoff 1991). They burrow head first into larger insect larvae. They hang to the surface film as pupae and may emerge in large numbers.

In 1999, the ceratopogonidae larvae were the second most abundant dipterans after the chironomids. They represented 189 of the 221 non-chironomid Diptera. All were of the genus *Bezzia* and *Probezzia*. The majority of the biting midges (111/189) were captured at location 8 in October.

Empididae

Empididae adults are known as "dance flies" and are small (<7mm) as larvae (Hilsenhoff 1991). The larvae are predaceous. The majority of Empididae larvae were recovered from location 7. In 1999, 11/12 and in 2001, 10/10 were recovered from this location. Only two were captured from location 12 in 2000. All were captured late in the summer (July 28 to August 28).

Dolichopodidae

These dipterans are commonly named, "long-legged flies." They are predaceous as both larvae and adults. Only one genus, *Rhaphium*, represented the 11 Dolichopodidae we captured. All of the *Rhaphium* were found at location 5 in 1999 and location 2 in 2001. Both of these sites could be considered to be fine sand beaches.

Coleoptera

The Coleoptera are commonly known as water beetles (Hilsenhoff 1991). In their larval phase they can swim, but commonly crawl along the bottom and respire cutaneously. As adults they have appendages adapted for swimming, eat with chewing mouthparts, and must surface to breathe. The requirement to surface generally restricts them to the shallow lake margins. A beetle generation takes one year of which approximately a month in spring is larval. They often use shoreline litter to pupate.

The use of substrate trays did not appear to be an effective way to capture aquatic beetles when compared to the benthic pump. In 1999, 14 beetles were recovered at five locations from 36 samples (Appendix 1, Table 2). Thirteen were from rocky sites and thirteen were larvae. In 2000, only four beetles were recovered from the 10 trays (Appendix 1, Table 3). The benthic pump in 2001 captured 96 beetles from 37 samples and a greater number of taxa (Appendix 1, Table 5).

Two genera of Dytiscidae dominated numerically in 2001, *Hydroporus* spp. (76/114) and *Brachyvatus* spp. (28/114). They are predaceous diving beetles as adults and range in size from 2-40 mm. The adults of these two genera were more abundant at night (64/65) where comparable day/night sampling was completed. The larvae of *Hydroporus* spp. were recovered from day samples collected at location 10.

Collembola

The adult Collembola (springtails) are considered to be semiaquatic, wingless arthropods, usually less than 3 mm long. They may reside on the surface of lotic waters and don't break through the surface film due to their small size and hydrophobic body surface. Thus, they are not benthic organisms. They specialize in items of food available on the water surface such as algae, detritus, and bacteria. The young springtails (nymphs) look similar to adults and we did not differentiate stages (Appendix 1, Tables 2-5).

Each year of sampling produced a different numerically dominant genus of springtail. In 1999, all 20 of the springtails recovered were *Anurida* sp. In 2000, 25 of 26 were *Isotoma* sp. In 2001, the majority (12/15) were *Sminthurus* sp.

Trichoptera

Trichoptera (caddisflies) construct portable cases out of plant material or sand. The larvae crawl along the bottom and are generally omnivorous or herbivorous. Most caddisflies pupate in their case. The adults feed on liquids and are attracted to lights at night (Clifford 1991; Wiggins et al. 1985).

Two families of Trichoptera dominated our catches during the three years of study. In 2001, Leptoceridae were the most numerically abundant (259/311) and were represented by two genera (*Mystacides* and *Nectopsyche*). Hydroptilidae were the next most abundant (41/311). In 2001, the majority of the caddisflies (309/311) were obtained from location 7 and location 2. In 1999, 24/50 caddisflies were obtained from location 7.

More caddisfly larvae were captured in August (240) than in May (43) from location 7 (Figure 21). However, this result was not significant (t-test, P = .09). All Hydroptilidae larvae were captured during the day, while more Leptoceridae larvae were captured at night (195) than during the day (47). However, this result was also not significant (Mann Whitney P=.18).

Heteroptera

The most numerous aquatic Heteroptera were the Corixidae. These "water boatmen" feed primarily on detritus, algae, protozoan, and very small invertebrates (Hilsenhoff 1991). The adults can fly and are common along the littoral zones of lakes. In 2001, fifty-six *Corisella* sp. adults were recovered. The majority (55/56) were recovered in May and August from location 7.

Hemiptera (Aphididae)

The aphids encountered during this study were of terrestrial origin. Although they were not identified to species, it is likely they were falling from riparian zone cottonwood trees (*Populus trichocarpa*) onto the water surface and onto our equipment and were thus included in our benthic samples. The majority of aphids (7/9 in 1999 and 25/28 in 2001) were recovered from locations 7 and 2. The riparian zones of these two locations consisted of stands of cottonwood trees.

A considerable number of aphids must have entered the lake foreshore from June to August. If we consider the total lake shore sampled, we estimated an aphid density of 1/1840 cm^2 in 1999 and 1/1550 cm^2 in 2001. These estimates equated to approximately 6 aphids per square meter of water surface. If we consider only the surface area sampled at the two cottonwood locations in 2001, than a density of 19 aphids per square meter of water surface can be estimated.

Ephemeroptera

The Insecta order Ephemeroptera are commonly called mayflies (Mounce 1973). The aquatic larvae occur among vegetation in littoral areas and sometimes along wave–swept beaches (Hilsenhoff 1991). The larva stage emerges from the water as a subimago and molts quickly into an adult. The adult and subimago stages of mayflies do not feed and are short-lived. Larval mayflies are mainly herbivore-detritivores.

The ecological characteristics and habitat preferences of the various families of Ephemeroptera identified in Shuswap Lake have been described by Thorp and Covich (1991), Mounce (1973) and Clifford (1991). The Baetidae are relatively small mayflies and are active swimmers, residing on the substratum. They tend to consume algae and detritus. Heptageniidae are relatively large and flat and are often found clinging to rocks. They feed on algae they scrape from the surface of rocks. Leptophlebiidae are medium sized invertebrates which and may be found within the gravel, in quiet waters. These mayflies were described as detritus feeders by Mounce (1973). Ephemeridae are relatively large filter-feeders that burrow into sand or silt, and may emerge in large numbers. Caenidae are small, can occupy bottom sediments, and emerge at the water surface. Surrounding vegetation and support is necessary for completion of their life cycle. Ephemerellidae may be found along wave-swept shores often within leaf litter. They crawl up on protruding substrate to emerge. Ameletidae (or Siphlonuridae) larvae are relatively large and mainly herbivores and detritivores.

In 1999, 311 mayfly nymphs were captured of which 50% were identified to species or genus (*Procloeon* sp. 104; *Ephemera simulans* 12; *Caenis simulans* 39). An average of 6.5/location (± 1.4 1SE) were counted. There was no significant difference between the numbers of mayfly nymph captured on cobble and sand/mud beaches (t-test, P = 0.11), although 226/311 mayflies were obtained from the rocky foreshore sites.

In 2000, Leptophlebiidae dominated and comprised 374 of 422 mayflies captured. An average of 42.4 mayflies/sample (± 6.3 1SE), were captured from the trays. Significantly more mayflies were captured in 2000 than in 1999 (Mann-Whitney Rank Sum P<.001). This may indicate they preferred the cleaner substrate used in 2000.

In 2001, 4 groups of mayflies dominated. Baetidae (603), Heptageneiidae (303), Caenidae (58), and Ameletidae (188) represented 89% of the 1289 mayflies recovered in 2001. *Ameletus* sp was most numerous in May, *Caenis simulans* and Heptageneiidae in June, and Baetids were common from June to late August (Appendix 1, Table 5).

DISCUSSION

The benthic species that dominated the foreshore from mid-May to mid-July varied each year, likely due to differences in sampling methods and differences in annual flooding characteristics. Cladocera, Chironomidae and Calanoida were the most numerous and most available to near-shore juvenile fish. Sididae were significantly more abundant in early spring. Oligochaeta and Nematoda were also abundant but occupied the interstitial spaces in the substrate and would not be readily available to juvenile salmonids. Ostracoda were abundant along the foreshore in May. However, the consumption of freshwater ostracods by juvenile salmon was not noted during preliminary analysis of Shuswap Lake chinook stomach samples (Brown and Winchell 2002). Ephemeroptera were common in 2000 and are

considered excellent fish food (Newman 2003). They are much larger than chironomids and although fewer in number, they represent an important item in the salmonids diet.

Many of B.C.'s lakes support rearing sockeye juveniles throughout the year (Stockner and Shortreed 1978; McDonald 1973). Although, we were sampling the benthic community while salmon fry appear to be surface oriented. It is possible that the large numbers of juvenile salmonids associated with the lake foreshore in spring may have influenced the number and type of benthic invertebrates we captured. In Quesnel Lake from mid-May until mid-June high densities of migrating sockeye fry were observed within 5 m of shore and these juvenile sockeye shifted their diet from *Leptodiaptomus* (near shore, water column) to pelagic Daphnia (limnetic) as the fish become more pelagic (Morton and Williams 1990).

The slightly higher numbers of Insecta noted on the lake foreshore at night can not be attributed to predator avoidance. The density of juvenile salmonids along the shallow lake margins was substantially higher in spring during the night than during the day (Brown and Winchell 2002). Thus, we might assume that nocturnal predation rates of invertebrates by fish would be higher at night than during the day. It is possible invertebrates may avoid shallow waters during the day to limit their exposure to UV light (Kelly and Bothwell 2002; Bothwell et al. 1994). The diurnal difference in invertebrate numbers was examined for only one sampling year (2001). There was no significant diurnal difference in total invertebrate numbers, or non-insecta numbers. The ostracods *Cypria* and *Cypris* were more numerous at night, while *Candona* were numerous during the day. Insecta were significantly more numerous at night and a significantly greater weight of insects was measured from the samples taken at night. Although slightly more Coleoptera, Trichoptera, Chironomidae, and Corixidae were noted from the night samples, there was no significant diurnal difference when each of the insect groups was examined separately.

There was no apparent difference in benthic communities between foreshore locations bordered by natural vegetation versus those areas cleared of trees. Our sampling design was not sophisticated enough to evaluate litter deposition rates or transport associated with lake wave actions. In the spring prior to maximum lake level, the unconsolidated foreshore separates the riparian vegetation from the water's edge. On beaches with a shallow gradient this distance can be over 100 meters. It is likely materials originating from the riparian zone are transported along the shore edge by wave action and are deposited upon certain shores and removed from others. Thus, the volume and type of litter that accumulates on the foreshore may not be related to the condition of the immediate uplands.

It did not appear that there was a relationship between the total number of invertebrates and substrate type. However, differences in the abundance of specific invertebrate families and species relative to habitat types were apparent. The cobble shores supported more Cladocera, especially the chydorids; *Acroperus harpae*, *Camptocerus rectorostris*, and *Alonella* spp. *Sida crystallina*, the most numerically abundant Cladocera, was also significantly associated with cobble sites. *Pleuroxus procurvatus* and *P. denticulatus* were associated with sand/mud substrate. More amphipods were captured from cobble/rocky foreshores. Slightly more chironomid larva were recovered from the sandy/mud foreshore locations in 1999 and 2001.

Location did appear to influence the number of invertebrates. We observed that certain locations produced greater numbers of invertebrates and supported differences in invertebrate community structure. Terrestrial aphids were consistently captured in greater numbers in benthic samples from two sites (location 7 and 2) where the upland areas were dominated by cottonwood trees. These two locations are also near the outlet of the Adams River and may be influenced by suspended materials washed down the river in the spring. Location 7 was associated with small beds of aquatic vegetation and was considered undisturbed. In 2001, location 7 supported a greater diversity of invertebrates and supported greater numbers of Oribatidae, Copepoda, Amphipoda, tubificids, chironomid larvae, Corixidae, Trichoptera, and bivalves.

ACKNOWLEDGMENTS

We would like to acknowledge the financial contribution of Fisheries Management in Kamloops (G. Kosikowski), Science Research (S. Macdonald and J. Pringle of MEHSD) and the ESSRF funds during 2000 and 2001. We especially appreciated the assistance given by H. Stalberg of Kamloops. Special consideration must be given to Applied Technical Services, Victoria. They sorted, identified, and counted each of the benthic invertebrates. We would like to thank B. Bravender and R. Russell for reviewing this manuscript.

INTERNET SOURCES

Fox, R. 2000. Biodiversity Lecture Outlines. Environmental Science 3000, Lader University. Freshwater Oligochaete Annelida. URL: http://www.lander.edu/rsfox/300oligoLec.html

Holmes, J.A. 2001. Brock University, Freshwater Ecology, BIOL 3P63, Lecture Notes. URL: http://www.utoronto.ca/env/lecture5.htm

Kramer, B. 2003. Shuswap Lake British Columbia. Peak Level History, Shuswap Lake Data Table, and Runoff Report 1999. URL: http://www.sicamous.com/users/kramerb

Newman, R. 2003. The Western Flyfisher . Insect Profiles. URL: http://bcadventure.com/adventure/angling/bugs/chironomid/chironomid.phtml

Nikon Microscopy. 2003. Digital Movie Gallery. Pond Life. URL: http://www.microscopyu.com/moviegallery/pondscum

Schuster, G. 2002. Freshwater Invertebrates, Biology 542/742. Gastropoda. URL: http://www.biology.eku.edu/SCHUSTER/bio%542/gastropoda.htm

Soil and Water Conservation Society of Metro Halifax (SWCSMH). 2002. Class Bivalvia, Oligochaetae, URL: http://lakes.chebucto.org/ZOOBENTH/BENTHOS/xxiv.html

Walter, D.E. 2002. Global Litter Invertebrate Decomposition Experiment (GLIDE). URL: www.nrel.colostate.edu/projects/glide/ostracoda.html

REFERENCES

Adams, M.A., Russell, D., and Haycock, D. 1990. Data report: Shuswap Lake monitoring program 1989. ECL Envirowest Consultants Limited. 204-800 McBride Boulevard, New Westminister, B.C.

Bothwell, M.L., Sherbot, D.M.J., and Pollock, C.M. 1994. Ecosystem response to solar ultraviolet-B radiation: Influences of trophic-level interactions. Science 265:97-100.

Brown, T.G. and Winchell, P. 2002. Use of Shuswap Lake foreshore by juvenile salmonids. A paper given at Institute of Ocean Sciences, Victoria, B.C. at MEHS All Staff meeting on Nov 27/2002. 11 p.

Chengalath, R. 1982. A faunistic and ecological survey of the littoral Cladocera of Canada. Can J. Zool. 60:2668-2682.

Clifford, H.F. 1991. Aquatic Invertebrates of Alberta. The University of Alberta Press. Edmonton, Alberta, Canada. 538 p.

Davies, M.E., Winsby, M.B., Gregory, R.S., and Levy, D.A. 1996. Effects of Eurasian watermilfoil on the ecology of Shuswap Lake. Hatfield Consultants Ltd. Report prepared for Department of Fisheries and Oceans, Canada. 40p + Appendices.

Delorme, L.D. 1991. Ostracoda, pp. 691-722. *In* Ecology and classification of North American freshwater invertebrates. Edited by J.H. Thorp and A.P. Covich. Academic Press, Inc., San Diego.

Dodson, S.I. and Frey, D.G. 1991. Cladocera and other Branchiopoda pp. 723-786. *In* Ecology and classification of North American freshwater invertebrates. Edited by J.H. Thorp and A.P. Covich. Academic Press, Inc., San Diego.

Edmondson, W.T. 1966. Fresh-Water Biology Second Edition. John Wiley and Sons, Inc. New York, USA. 1248 p.

Fedorenko, A.Y. and Pearce, B.C. 1982. Trapping and coded wire tagging of wild juvenile chinook salmon in the South Thompson/Shuswap River System 1976, 1979, 1980. Can. Manuscr. Rep. Fish. Aquat. Sci. 1677: 63 p.

Graham, C.C. and Russell, L.R. 1979. An investigation of juvenile salmonid utilization of the delta-lakefront area of the Adams River, Shuswap Lake. Can. Fish. Mar. Serv. Manuscr. Rep. 1508: 32 p.

Hatfield Consultants Ltd. 1996. Effects of Eurasian watermilfoil on the ecology of Shuswap Lake. Report prepared for C. Levings, Department of Fisheries and Oceans, Canada, by Hatfield Consultants Ltd. 201-1571 Bellevue Ave. West Vancouver, B.C. 97 p + Appendices.

Hilsenhoff, W.L. 1991. Diversity and classification of insects and Colembola, pp. 593-664. *In* Ecology and classification of North American freshwater invertebrates. Edited by J.H. Thorp and A.P. Covich. Academic Press, Inc., San Diego.

Kelly, D.J. and Bothwell, M.L. 2002. Effects of solar radiation on shallow stream invertebrates. National Institute of Water and Atmospheric Research, Christchurch, New Zealand. 3 p.

McDonald, J. 1973. Diel vertical movements and feeding habits of underyearling sockeye salmon (*Oncorhynchus nerka*), at Babine Lake, B.C. Fish. Res. Bd. Can. Tech. Rep.

Morton, K.F. and Williams, I.V. 1990. Sockeye salmon (*Oncorhynchus nerka*) utilization of Quesnel Lake, British Columbia. Can. Tech. Rep. Fish. Aquat. Sci. 1756: 29 p.

Mounce, D.E. 1973. An introductory guide to stream insects of Southern Vancouver Island. Fisheries Research Board of Canada. Pacific Biological Station, Nanaimo, B.C. Circular No. 95. 39 p.

Russell, L.R., Graham, C.C., Sewid, A.G., and Archibald, D.M. 1980. Distribution of juvenile chinook, coho and sockeye salmon in Shuswap Lake – 1978-1979; Biophysical inventory of littoral areas of Shuswap Lake, 1978. Can. Fish. Mar. Serv. Manuscr. Rep. 1479: 54 p.

Smith I.M. and Cook, D.R. 1991. Water mites, pp. 523-592. *In* Ecology and classification of North American freshwater invertebrates. Edited by J.H. Thorp and A.P. Covich. Academic Press, Inc., San Diego.

Southwood, T.R.E. 1966. Ecological Methods, with Particular Reference to the Study of Insect Populations. Methuen & Co. Ltd., London.

Stockner, J.G. and Shortreed, K.R.S. 1978. Limnological survey of 35 sockeye salmon (*Oncorhynchus nerka*) nursery lakes in British Columbia and the Yukon Territory. Can. Dep. Fish. Envir., Fish. Mar. Serv., Tech. Rep., 827: 47 pp.

Thorp, J.H. and Covich, A.P. 1991. Ecology and classification of North American freshwater invertebrates. Academic Press, Inc. New York, USA. 911 p.

Wiggins, G.B., Weaver III, J. S. and Unzicker, J.D. 1985. Revision of the caddisfly family Uenoidae (*Trichoptera*). The Canadian Entomologist 117:763-800.

Williams, I.V., Gilhousen, P., Saito, W., Gjernes, T., Morton, K., Johnson, R., and Brock, D. 1989. Studies of the lacustrine biology of the sockeye salmon (*Oncorhynchus nerka*) in the Shuswap system. International Pacific Salmon Fisheries Commission. Bulletin XXIV 108 p.

Williamson, C.E. 1991. Copepoda, pp. 787-822. *In* Ecology and classification of North American freshwater invertebrates. Edited by J.H. Thorp and A.P. Covich. Academic Press, Inc., San Diego.

LIST OF FIGURES

1. Extent of Shuswap Lake foreshore in early spring prior to rise in lake level. Photo taken looking towards Location 2.

2. A cobble beach (Location 18).

3. Main arm of Shuswap Lake, illustrating the benthic sampling locations for three years of study.

4. Plastic benthic trays used in 1999-2000.

5. Placement of cobble and rock into rows at location 12.

6. Benthic pump used in 2001

7. Relative benthic invertebrate composition of benthic substrate trays collected in spring 1999.

8. Relative benthic invertebrate composition of benthic substrate trays collected in spring 2000. Substrate consisted of washed and sorted fine gravel.

9. Relative benthic invertebrate composition of samples collected in spring and summer 2001 with a benthic pump sampler.

10. Number of benthic invertebrates for each of three springs (1999-2001).

11. Relative composition of benthic samples for each of three springs (1999-2001). Major invertebrate groups are illustrated and differences between their relative numeric compositions are likely due to different sampling methods.

12. The relationship between the average volume of sediment (3 samples per location) and average number of organisms. Each sample represents 1725 cm^3 of substrate. Cobble/rock and sand/mud locations are indicated by data-labels.

13. Relationship between the average volume of sediment at each location (3 samples per location) and average number of non-insecta organisms. A sample represents 1725 cm^3 of substrate. Location numbers are indicated by data-labels.

14. Total number of benthic invertebrates (sum of 3 samples) at each of 16 locations in 1999. Substrate (cobble or sand/mud) and status of riparian fringe (cleared or treed) is indicated.

15. Total number of benthic invertebrates per sample taken in 2000.

16. Numbers of Hydracarina, Ostracoda, Copepoda, and three families of Cladocera per benthic pump sample. Foreshore samples were taken in May and August 2001 at location 7 during day and night.

17. Numbers of bivalves, amphipods, nematodes, and Oligochaeta (Tubificidae, Enchytraeidae, and Naididae) per benthic pump sample. Foreshore samples taken in May and August 2001 at location 7 during both day and night.

18. Numbers of 3 Oligochaeta families from all samples taken in 2001.

19. Relative numbers of Insecta from all samples taken in 2001.

20. Relative weight of Insecta from all samples taken in 2001.

21. Numbers of Insecta (Orders) per benthic pump sample. Foreshore samples were taken in May and August 2001 at location 7 during both day and night.

22. Weight of insects / benthic pump sample. Foreshore samples taken during both day and night. Location 7 is represented by samples 1-6 (May) and 25-30 (August).

23. Density of chironomid larvae and pupae captured during each of three years.

24. Number of chironomid larvae/benthic pump sample. Foreshore samples taken during both day and night. Location 7 is represented by samples 1-6 (May) and 15-30 (August).

FIGURES

Figure 1. Extent of Shuswap Lake foreshore in early spring prior to rise in lake level. Photo was taken looking towards Location 2.

Figure 2. A cobble beach (location 18).



Figure 3. Main Arm of Shuswap Lake, illustrating the benthic sampling locations for three years of study.



Figure 4. Plastic benthic trays used in 1999-2000.

Figure 5. Placement of cobble and rock into rows at location 12.



Figure 6. Benthic pump used in 2001.



Figure 7. Relative benthic invertebrate composition of benthic substrate trays collected in spring 1999.



Figure 8. Relative benthic invertebrate composition of benthic substrate trays collected in spring 2000. Substrate consisted of washed and sorted fine gravel.



Figure 9. Relative benthic invertebrate composition of samples collected in spring and summer 2001 with a benthic pump sampler.



Figure 10. Number of benthic invertebrates for each of three springs (1999-2001).



Figure 11. Relative composition of benthic samples for each of three springs (1999-2001). Major invertebrate groups are illustrated and differences between their relative numeric compositions are likely due to different sampling methods.



Figure 12. The relationship between the average volume of sediment (3 samples per location) and average number of organisms. Each sample represents 1725 cm^3 of substrate. Cobble/rock and sandy foreshore locations are indicated by data-labels.



Figure 13. Relationship between the average volume of sediment at each location (3 samples per location) and average number of non-insecta organisms. A sample represents 1725 cm^3 of substrate. Location numbers are indicated by data-labels.



Figure 14. Total number of benthic invertebrates (sum of 3 samples) at each of 16 locations in 1999. Substrate (cobble or sand/mud) and status of riparian fringe (cleared or treed) is indicated.



Figure 15. Total number of benthic invertebrates per tray sample . Samples were taken on sandy/mud or cobble/rock foreshore. Control samples consisted of bucket hauls with empty trays (indicates invertebrates in water column).



Figure 16. Numbers of Hydracarina, Copepoda, Ostracoda, and Cladocera per benthic pump sample. Foreshore samples were taken in May and August 2001 at location 7 during both day and night.



Figure 17. Numbers of bivalves, amphipods, nematodes, and Oligochaeta (Tubificidae, Enchytraeidae, and Naididae) per benthic pump sample. Foreshore samples taken in May and August 2001 at location 7 during both day and night.



Figure 18. Numbers of 3 Oligochaeta families from all samples taken in 2001.



Figure 19. Relative numbers of Insecta from all samples taken in 2001.



Figure 20. Relative weight of Insecta from all samples taken in 2001.



Figure 21. Number of Insecta (Orders) per benthic pump sample. Foreshore samples taken in May and August 2001 at location 7 during both day and night.



Figure 22. Weight of insects / benthic pump sample. Foreshore samples taken during both day and night. Location 7 is represented by samples 1-6 (May) and 25-30 (August).



Figure 23. Density of chironomid larvae and pupae captured during each of three years with 95% confidence limits (CL).



Figure 24. Number of chironomid larvae/benthic pump sample. Foreshore samples taken during both day and night. Location 7 is represented by samples 1-6 (May) and 15-30 (August).

APPENDIX 1. BENTHIC SPECIES LIST; SHUSWAP LAKE FORESHORE

Group	Family	Genus
COELENTERATA		
		Hydra
TURBELLARIA		
NEMATODA		
BRYOZOA		
OLIGOCHAETA		
	Enchytraeidae	
	Naididae	
		Chaetogaster
		Nais
		Pristina foreli
		Stylaria lacustris
		Vejdovskyella
		Slavinia appendicularia
	Tubificidae	
	Lumbriculidae	
	Lumbricidae	
HIRUDINEA		
		Piscicola salmositica
		Dina parva
		Helobdella stagnalis
TARDIGRADA		
ARACHNIDA		
HYDRACHNIDIA(Hydracarina)		
		Hydracarina #1
		Frontipoda
		Torrenticola
		Forellia
	Oribatidae	
		Hydrozetes
OSTRACODA		
		Candona
		Cyclocypris
		Limnocythere
		Cypria
		Cypris
		Cypricercus elongata
COPEPODA		
Calanoida		
		Cyclops scutifer
		Eucyclops agilis
		Leptodiaptomus
		Epsichura nevadensis
		Macrocycops ater
Cyclopoida		
		Cyclops capillatus
Harpacticoida		
	Canthocamptidae	
CLADOCERA		
	Daphnidae	
		Daphnia
		Ceriodaphnia

Scapholeberis mucronata
Scapholeberis rammneri

Leptodoridae

Leptodora kindti

Macrothricidae

Ilyocryptus spinifer

Chydoridae

Acroperus harpae
Alona costata
Alona guttata
Alonella excisa
Camptocercus rectorostris
Chydorus sphaericus
Eurycercus lamellatus
Pleuroxus denticulatus
Pleuroxus procurvatus
Graptoleberis testudinaria

Sididae

Sida crystallina

Polyphemidae

Polyphemus pediculus

AMPHIPODA

Hyalella azteca

INSECTA
DIPTERA

Chironomidae

Tanypodinae

Paramerina
Procladius
Thiennemannimyia

Chironominae

Tanytarsini

Micropsectra
Cladotanytarsus
Paratanytarsus
Rheotanytarsus
Stempellina
Tanytarsus
Stempellinella

Chironomini

Chironomus
Demicryptochironomus
Paratendipes
Microtendipes
Paracladopelma
Stictochironomus
Phaenopsectra
Polypedilum (pentapedilum)

Orthocladiinae

Corynoneura
Cardiocladius
Cricotopus
Epoicocladius
Orthocladius
Eukiefferiella

Heterotrissocladius
Paracladius
Psectrocladius
Thienemanniella

Diamesinae

Potthastia

Podonomidae

Trichotanypus

Ceratopogonidae

Bezzia
Probezzia

Muscidae

Limnophora

Tabanidae

Tabanus

Cecidomyidae
Empididae

Phyllodromia

Phoridae

Megaselia

Sciomyzidae
Dolichopodidae

Rhaphium

HYMENOPTERA

Chalcoidea
Formicidae
Tenthredinidae
Encyrtidae

HOMOPTERA

Aphididae

COLEOPTERA

Haliplidae

Haliplus

Dytiscidae

Brachyvatus
Hydroporus
Oreodytes

Curculionidae
Staphylinidae
Phalactidae
Gyrinidae

Gyrinus

LEPIDOPTERA
EPHEMEROPTERA

Baetidae

Baetis tricaudatus
Procloeon

Heptageneiidae

Heptageneia
Cinygmula
Heptagenia solitaria

Leptophlebiidae

Choroterpes albiannulata
Paraleptophlebia bicornuta
Paraleptophlebia temporalis

	Leptophlebia
Ephemeridae	
	Ephemera simulans
	Hexagenia limbata
Caenidae	
	Caenis simulans
Ephemerellidae	
	Ephemerella inermis
Tricorythidae	
	Tricorythodes minutus
Ameletidae	
	Ameletus
PLECOPTERA	
Chloroperlidae	
Perlodidae	
	Suwallia
TRICHOPTERA	
Hydroptilidae	
	Agraylea
Lepidostomatidae	
	Hydroptila
	Lepidostoma
Leptoceridae	
	Mystacides
	Nectopsyche
Brachycentridae	
Limnephilidae	
	Onocosmoecus
	Brachycentrus
HEMIPTERA	
Corixidae	
	Corisella
THYSANOPTERA	
COLLEMBOLA	
Hypogasturidae	
	Anurida
Isotomidae	
	Isotoma
Sminthuridae	
	Sminthurus
ODONATA	
Anisoptera	
Aeschnidae	
	Aeschna umbrata
Coenagridae	
Zygoptera	
	Enallagma sp.
	Enallagma cythigerum
MOLLUSCA	
Gastropoda	
Planorbidae	
	Gyraulus
Lymnaeidae	
	Stagnicola
	Lymnaea
Physidae	

Physa

Valvatidae

Valvata sincera sincera

Pelecypoda

Sphaeriidae

Sphaerium
Pisidium

APPENDIX 2. SHUSWAP LAKE BENTHIC SAMPLES (1999-2001)

Table 1. Number of non-Insecta benthic invertebrates from Shuswap Lake foreshore in 1999.

Table 2. Number of benthic Insecta from Shuswap Lake foreshore in 1999.

Table 3. Number of benthic invertebrates from Shuswap Lake foreshore in 2000.

Table 4. Number of non-Insecta benthic invertebrates from Shuswap Lake foreshore in 2001.

Table 5. Number of benthic Insecta from Shuswap Lake foreshore in 2001.

Table 6. Weight (mg) of benthic Insecta from Shuswap Lake foreshore in 2001.

Table 1. Number of Non-Insecta Benthic Invertebrates from Shuswap Lake Foreshore in 1999

Location		12			17			18			16			13			15			6			14			10			5			9			4			2			1			7		8				
Date		July 28			July 28			July 29			July 29			July 29			July 29			July 29			July 29			July 30			July 30			July 30			July 30			July 30			July 30			July 30		Oct 04				
Replicate		1	2	3	1	2	3	1	2	3	1	2	3	1	2	3	1	2	3	1	2	3	1	2	3	1	2	3	1	2	3	1	2	3	1	2	3	1	2	3	1	2	3	1	2	3		
Benthos #		1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	
Sample #		514	515	516	517	518	519	520	521	522	523	524	525	526	527	528	529	530	531	532	533	534	535	536	537	538	539	540	541	542	543	544	545	546	547	548	549	550	551	552	553	554	555	556	557	558	559	560	561	TOTAL	SUM
Species	Stage																																															
COELENTERA																																																	
Hydra									1						1	1				2			1																											
Turbellaria												11		24	1						1				2	3	11						1		4													
Nematoda		1	3	1	70	50	50	21	2	71	298	74	188	72	40	41	1		1	13	1	96		1	4	51	82	43	18		3		5		20	24	55	76	31	19	183	120	94	56	138	17	162	91	443	
Tardigrada											10				10								1							2	5				1					1				10		10	10			
Bryozoa						1																																											
Oligochaeta	egg																									3																								
Oligochaeta	juv	1	4	4	100	90	60	120	60	90	150	60	60	90	160	90	2	14	9	180	50	150	9	37	32	100	60	20	47	29	130	50	50	65	70	92	40	180	30	5	70	50		100	70	70	566	464	1790	
Naididae		20	18	34	50	27	7	7	25	68	19	26	25	9	7	4		1	5	32	11	30	1	2	3	5	18	30			5	1		1	3	1		7	4			7	1	34	24	15	14		9	
Chaetogaster		5	3	3	48	36	12	3	14	25	2	5	16	43	33	33			2	8	4	9	6	11	4	3	12		21	16	80	15	22	56	40	29	12	10	1	3	26	10		18	1	10		121	143	
Nais				2	32	17	8	11	19	27	21	158	55			2	1	1	1	31	10	3	1			2	1			1	2	2	4	8	7	7	1	58		3	5	5	1	20	198	130	12	1	28	
Stylaria lacustris				3	31	55	61	11	15	14	3		1		2	2				1		1				2	2	1			1	1	2	1				1	1	1		1			7	3					
Pristina foreli								7	4	4	46	65	50		1	1				37	60	23		2	4	3	13	21	9	13	16	14	46	90	5	3	2	93	10	5	90	50	3	1			11	4	39	
Slavinia appendicularia								1	1		3		1	4						4	1	2																							2	1				
Tubificidae												1																																		15	34	55	
Lumbriculidae				1																	4						1																						
Lumbricidae				1																					1																								
Enchytraeidae				2	1			10	2	4	5		1		1					3	8	6			3	11	60	7	2	1	1			1	6	2	10		1	1	12		22	3	7	12	2	29	
Hirudinea																																																	
Piscicola salmonitica																					1																												
Hydracarina				1	8	16	20	3	16	22				29	12	12		1		33	2	1		1		5	14	23				1	30	4	6		1	4	1			10							
Hydracarina #1	(parasitic)				10			10	10					10																																		
Orbatidae												10								20				1		20					2			5	1	2		10		10			60	10	10				
Hydrozoles				1	11			10	10	20		10		20					1					3			10	2	1	1					1										10	10			1	
Ostracoda									20	10																													5										
Candona				1				10						10						60	240	110				51	10	60												1			1	1	21	1	
Cyclocypris				1																																										1		
Cypria									20		10		1														1														10			1		6	
Cypris								31		1				11						11	10														1				1								
Calanoida	nauplius						10				20				10								1	4	1		10		3	4	1	10	40	10	20	5	6	20	15	1							23		
Calanoida	copepodite																							1																									
Cyclops scutifer																													1																			
Eucyclops agilis		1			6	34	28	45	8	45	163	63	236	42	32	16	1	3	5	27	50	7		20	6	2	21	60	9	12	30	5	8	8	5	13		245	65	23	90	10	4	3	23	39		42	63	
Leptodiaptomus					1						1	1	2		10	1				10	1	1		1	3				131	113	135	5			15	10	8	1	12	4	10		1	10	2	1			
Epischura											1		1	2	1			1					37	9	16	4	3	1																				
Macrocyclops albidus					3				20	33	7	1	3	3	2	5		6		18	15	10		1		4	1			1		4	1	4			1	1		5			1	3				
Cyclopoida	copepodite		2		170	71	140	240	315	312	165	92	208	111	218	152	1	18	12	134	210	128	5	96	17	102	131	350	26	33	35	70	165	70	40	20	12	610	85	17	90		80	141	258	202	11	267	169	
Cyclopoida	nauplius				10					10			10		30	10		5	2	90	140	40		5		20	50	20	3			10	120	20		6	4	30	15	1	30	70	10		40			40	
Harpacticoida	copepodite																													3																		
Harpacticoida																													1																		
Harpacticoida	nauplius						20																						2	2		60	60	25		3	2	230	45	14	110	70	10	10	10	30		60	20	
Canthocamptidae					20			10											1	30	30	1							4	2	5				2					20	20				10		100	81	
Cladocera		1																																														
Daphnidae																																																
Daphnia														1				1		2	1		1		3		1											1									
Ceriodaphnia																																								5		2						
Scapholeberis mucronata						10																				10												10	3	1			14	32	7				
Macrothricidae																																																
Ilyocryptus spinifer																																																
Chydoridae								70	60	200	360	70	140	30	30	50	2			270	460	310	3	18	8	100	220	100	82	43	50	110	20	5	80	21	4	310	60	3		40		70	150	90		20	20	
Acroperus harpae				11	42	20	71	16	212		10		13	31	14		1	1		295	80	74	1	20	1	53	81	50	1		1		110	40		1		6		15	10			33	1				
Alona (costata?)		2	1	50	18	30	20	22	2		80	11	10	32	10					56	10	102				14	32	20	1	1	1				10	1		8	6	1	53	11			12		20	1	
Alona sp. 2				20	10	70	10	30	171	141	40	100	10	10	40		6	4		160	280	190	1	4	4	140	120	80	46	75	100		10	15	35	15	5	100	30	1		20			20			60	
Camptocercus rectirostris													21							30				3		3	21	20					217	165		1				1		25	82	32				
Chydorus sphaericus			1		40	70	70	93	104	96	131	90	190	100	21	101		3		252	600	121	3	10	5		60		10	7	5	25	60	25	3	2	2	51	16	7	40	20		31	483	55		100	60	
Eurycercus lamellatus							1							2	7	6				10		10		1												1	2	1										
Pleuroxus denticulatus																								1		20				2								30		17	31	20							
Alonella																				30	190	230		4	1		100	60	19							9	3	23											
Pleuroxus procurvatus																																						20		1	10			20	1				
Sididae																																																	
Sida crystallina			12	15	45	53	69	9	40	46	5	2	4	15	36	75	1	5		284	83	67		11	3	41	109	3			1	298	243	761	7	1	4			1		15	67	16				
Polyphemidae																																																	
Polyphemus pediculus																																												1	10	1				
Amphipoda																						5																											
Hyallela azteca					2	6	4	12	35	37		2	1	17	21	7					13					17	11	11				2							1					2	5	5			7	
Gastropoda																						3		1																								1	
Stagnicola														1					1	1																													
Physa						1		3	1	4				1								1				1																						
Gyraulus																																															1	
Bivalvia																																																
Sphaerium																																															1	
Fish																																																
Cottus asper	1																			1																												
TOTAL		31	44	70	739	599	690	828	879	1564	1580	911	1494	721	779	693	9	68	49	2135	2585	1772	74	276	119	791	1201	1046	442	364	607	684	1214	1375	373	279	166	2106	470	113	878	568	236	643	1668	811	816	1370	3069	

Table 1. Number of Non-Insecta Benthic Invertebrates from Shuswap Lake Foreshore in 1999

Location		12			17			18			16			13			15			6			14			10			5			9			4			2			1			7			8		
Date		July 28			July 28			July 29			July 29			July 29			July 29			July 29			July 29			July 30			July 30			July 30			July 30			July 30			July 30			July 30			Oct 04		
Replicate		1	2	3	1	2	3	1	2	3	1	2	3	1	2	3	1	2	3	1	2	3	1	2	3	1	2	3	1	2	3	1	2	3	1	2	3	1	2	3	1	2	3	1	2	3	1	2	3
Benthos #		1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48
Sample #		514	515	516	517	518	519	520	521	522	523	524	525	526	527	528	529	530	531	532	533	534	535	536	537	538	539	540	541	542	543	544	545	546	547	548	549	550	551	552	553	554	555	556	557	558	559	560	561
Species	Stage																																															
COELENTERA																																																
Hydra									1					1	3					2			1																										
Turbellaria												11		24	1							1				2	3	11							4												
Nematoda		1	3	1	70	50	50	21	2	71	298	74	188	72	40	41	1		1	13	1	96		1	4	51	82	43	18		3		5		20	24	55	76	31	19	180	120	94	56	128	17	162	91	443
Tardigrada											10				10								1							2	5					1				1			10			10	10		
Bryozoa						1																																										
Oligochaeta	egg																									3																							
Oligochaeta	juv	1	4	4	100	90	60	120	60	90	150	60	60	90	160	90	2	14	9	180	50	150	9	37	32	100	60	20	47	29	130	50	50	65	70	92	40	180	30	5	70	50		100	70	70	566	464	1790
Naididae		20	18	34	50	27	7	7	25	68	19	26	25	9	7	4		1	5	32	11	30	1	2	3	5	10	30			5	1		1	3	1		7	4			7	1	34	24	15	14		9
Chaetogaster		5	3	3	48	36	12	3	14	25	2	5	16	43	33	33			2	8	4	9	6	11	4	3	12		21	16	80	15	22	56	40	29	12	10	1	3	26	10		18	1	10		121	143
Nais				2	32	17	8	11	19	27	21	158	55			2	1	1	1	31	10	3	1			2	3			1	2	2	4	8	7	7	1	58		3	5	5	1	20	198	130	12	1	28
Stylaria lacustris				3	31	55	61	11	15	14	3		1		2	2				1		1				2	2	1			1	1	2	1				1	1	1		1			7	3			
Pristina foreli								7	4	4	46	65	50		1	1				37	60	23		2	4	3	13	21	9	13	16	14	46	90	5	3	2	93	10	5	90	50	3	1			11	4	39
Slavina appendiculata								1	1		3		1	4						4	1	2																							2	1			
Tubificidae												1																																		15	34	55
Lumbriculidae					1																4						1																					
Lumbricidae					1																				1																				1			
Enchytraeidae					2	1			10	2	4	5		1		1				3	8	6				3	11	60	7	2	1	1			1	6	2	10		1	1	12		22	3	7	12	2	29
Hirudinea																						1																										
Piscicola salmositica																																															
Hydracarina				1	8	16	20	3	16	22				29	12	12		1		33	2	1		1		5	14	23				1	30	4	6		1	4	1			10						1
Hydracarina #1	parasites				10			10	10					10																						5	1	2		10		10			60	10	10		
Oribatidae												10									20			1		20			2																		
Hydrachna				1	11			10	10	20		10		20					1					3			10	2	1	1							1								10	10			1
Ostracoda									20	10																													5									
Candona				1				10						10						60	240	110				51	10	60													1				1	1	21	1
Cyclocypris				1																							1																1		1		6
Cypria									20		10		1																												10						
Cypris								31		1				11						11	10						1									1										23		
Calanoida	nauplius						10				20				10								1	4	1		10		3	4	1	10	40	10	20	5	6	20	15	1								
Calanoida	copepodite																							1																								
Cyclops scutifer																													1																		
Eucyclops agilis			1		6	34	28	45	8	45	163	63	236	42	32	16	1	3	5	27	50	7		20	6	2	21	60	9	12	30	5	8	8	5	13		245	65	23	90	10	4	3	23	39		42	63
Tropodiaptomus					1							1	2		10	1				10	1	1		1	3			131	113	135	5				15	10	8	1	12	4	10		1	10	2	1		
Epischura											1		1	2	1		1						37	9	16	4	3	1			1																
Macrocyclops ater					3				20	33	7	1	3	3	2	5		6		18	15	10		1		4	1			1		4	1	4				1	1		5			1	3			
Cyclopoida	copepodite		2		170	71	140	240	313	312	165	92	208	111	218	152	1	18	12	134	210	128	5	96	17	102	131	350	26	33	35	70	165	70	40	20	12	610	85	17	90		80	141	258	202	11	267	169
Cyclopoida	nauplius				10					10			10		30	10		5	2	90	140	40		5		20	50	20	3			10	120	20		6	4	30	15	1	30	70	10		40			40
Harpacticoida	copepodite																												1	3																	
Harpacticoida																													1																	
Harpacticoida	nauplius						20			40	20	40	190	20	20	20		2	4	20	20	40	3	8	3	10			2	2		60	60	25		3	2	230	45	14	110	70	10	10	10	30		60	20
Canthocamptidae					20		10												1	30	30	1							4	2	5				2					20	20				10		100	81
Cladocera			1																																												
Daphnidae																																							1									
Daphnia														1				1		2	1		1		3		1													5		2						
Ceriodaphnia																																															
Scapholeberis mucronata						10																				10											10	3	1			14	32	7				
Macrothricidae																																															
Ilyocryptus spinifer																																	1														
Chydoridae								70	60	200	360	70	140	30	30	50	2			270	460	310	3	18	8	100	220	100	82	43	50	110	20	5	80	21	4	310	60	3		40		70	150	90		20	20
Acroperus harpae					11	42	20	71	16	212		10		13	31	14		1	1	295	80	74	1	20	1	53	81	50	1		1		110	40		1		6		15	10			33	1			
Alona (costata?)		2	1		50	10	30	20	22	2		80	11	10	32	10				56	10	102				14	32	20	1	1	1				10	1		8	6	1	53	11			12		20	1
Alona sp. 2					20	10	70	10	30	171	141	40	100	10	10	40		6	4	160	260	190	1	4	4	140	120	80	46	75	100		10	15	35	15	5	100	30	1		20				20		60
Camptocercus rectirostris														21						30				3		21	20					217	165		1					1								
Chydorus sphaericus			1		40	70	70	93	104	96	131	90	190	100	21	101		3		252	600	121	3	10	5		60		10	7	5	25	60	25	3	2	2	51	16	7	40	20		31	483	55		100	60
Eurycercus lamellatus							1							2	7	6				10		10		1												3	2	1									
Pleuroxus denticulatus																								1		20				2								20		17	31	20						
Alonella																				30	190	230		4	1		100	60	19						9	3	23											
Pleuroxus procurvatus																																					20		1	10				20	1			
Sididae																																															
Sida crystallina			12	15	45	53	69	9	49	46	5	2	4	15	36	75	1	5		284	83	67		11	3	41	109	3			1	298	243	761	7	1	4				1		15	67	16			
Polyphemidae																																															
Polyphemus pediculus																						5																					1	10	1				
Amphipoda																					33					17	11	11			2							1				2	5	5			7	
Hyalella azteca					2	6	4	12	35	17		2	1	17	21	7						3		1																							1	
Gastropoda														1					1	1																												
Stagnicola																						1																										
Physa						1		3	1	4				1												1																						1	
Gyraulus																																															
Bivalve																																															
Sphaerium																																															
Fish																																															
Coitus asper	1																			1																												
TOTAL		31	44	70	739	599	690	828	879	1564	1580	911	1494	721	779	693	9	68	49	2135	2585	1772	74	276	119	791	1201	1046	442	364	607	684	1214	1375	373	279	166	2106	470	113	878	568	236	641	1668	811	816	1370	3069

Table 2. Number of Benthic Insecta from Shuswap Lake Foreshore in 1999

Location		12			17			18			16			13			15			6			14			10			5			9			4			2			1			7		8		
Date		July 28			July 28			July 29			July 29			July 29			July 29			July 29			July 29			July 30			July 30			July 30			July 30			July 30			July 30			July 30		Oct 04		
Replicate		1	2	3	1	2	3	1	2	3	1	2	3	1	2	3	1	2	3	1	2	3	1	2	3	1	2	3	1	2	3	1	2	3	1	2	3	1	2	3	1	2	3	1	2	3
Benthos #		1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48
Sample #		514	515	516	517	518	519	520	521	522	523	524	525	526	527	528	529	530	531	532	533	534	535	536	537	538	539	540	541	542	543	544	545	546	547	548	549	550	551	552	553	554	555	556	557	558	559	560	561
Species	Stage																																														
Insecta																																															
Chironomidae	L	17	18	8	233	128	58	262	190	727	100	10	41	322	466	384	2	2	2	145	70	94		15	2	40	140	20	26		15	15	120	175	5	14	4	63	15	9		1	21	136	43	245	672	978
Chironomidae	P														2																																	
Tanypodinae	L										10			25	67	6		1		2	1	5	1	2	1	23	2	5	3			1	5	1				5	2	7		1	29	12	15	1	1	7
Paramerina	L			1										10	7	8				2	2	3				1	5	1			1	1	3	1	1								1				
Paramerina	P													3		1								1		2	1					1															
Procladius	L							1																																1			1		1	39	25
Thienemannimyia	P				1																																									
Thienemannimyia	L	1	6	3	5	6	9	2	4	8				5	4	2	1			2	1	6		1	3	6	7	6		1									5		3			1				
Tanytarsini	L	5	10	7	53	31	23	40	40	96	22	13	81	126	251	152	1			156	50	150	1	3	1	93	98	20	17	5	12	16	46	17	12	6	2	120	7	16		34	10	68	120	257	606	1994
Tanytarsini	P																					1																										
Micropsectra	L			1	1		1	1	4	5				2	1				1		5	9			1	5	1				1							4			1	12	3	3	1			
Parastenpellina	L		5	1	2				4	2	10	10	29	7	13	2				8	5	26						1	19	1	7				7	3	1	17		3	9	8	2	38	31	5	4	19
Parastenpellina	P											1																																			
Stempellina	P							1							1											1																						
Stempellina	L		1	1					1	3					1						2					4	1	1				1									1						
Tanytarsus	L	5	18	18	2	1	1	2	15	16	4	6	3	27	31	8				10	2	31				2	6	2				2		2	2		1	4	1	5		12	14	25	25	2		5
Tanytarsus	P	1		2		2				1										3	2			1			1				1											1					
Stempellinella	P																																													42
Chironomini	L							22	4	1			10	43	12	30	1			2	1	3				13			3						9	5	6	2		1		3	3	12	287	286	420
Chironomini	P																																														
Chironomus	P																																								1	1						
Chironomus	L	1								1	1	1																													3	2		3	2	8			
Demicryptochironomus	L			1	2		1		2	1		2	1	7	5	4				6	1	4				3	1	3	2	3	3			4	2	2	1	1		1	3	1	2	16	13	51
Demicryptochironomus	P																																														
Parastendipes	P							1						1														1																			
Parastendipes	L				2	1		23	11	3				9		1				2		3					2	2														1					36
Phaenopsectra	L																					1				4															4	7	5	6	5	8
Polypedilum (pentapedilum)	L																																											279	63	164
Polypedilum (polypedilum)	L																												7	1													1	1		1
Orthocladiinae	P																1															1													1	
Orthocladiinae	L		1	2	11	16	45	54	56	118	11	2	3	40	81	170				43	17	36	2	7		10	1		2		1	1	26	11	2	3		8	3		2	2	3	46	5	1	62	37
Corynoneura	L	1	1	2	268	217	155	129	76	168	113	20	59	123	107	361	2			99	112	182		64	24	89	193	140	29	9	36	50	29	36	41	14	1	85	25	11	4	13	11	76	30			1
Corynoneura	P				2	1		1		1										22	1						13						1							1	1						
Cricotopus	L																			1		3																									
Eukiefferiella	L																												2																		
Heterotrissocladius	L		1																		1	3																			1						
Orthocladius	L																																	1							1		4	1			
Paracladius	L				7	1		1		1						2																															
Psectrocladius	P		1																																												
Psectrocladius	L			1	7	6	1	4	8	8		2	1	4	4	19				2									3	1	2	1	2	1							3						
Thienemanniella	L		1			2		1	1	4					11					23	12	78	16			1	10						5	1	1						1	22	8				1
Tanypodinae	L																																														
Potthastia	L																																											4			
Diptera																				10																												
Ceratopogonidae							10							20																		1	10											26	19	50
Bezzia	L				1																											11			5	5	1	5			2					
Probezzia	L				1						1				1		1	1	1							1																	9	2	5
Empididae	L									1																															10		1				
Phyllodromia	L																																			1	1	1	1								
Pleuridae	L																																							1							
Megaselia	L																														1															
Sciomyzidae	P													1																																
Dolichopodidae	L																																														
Rhaphium	L																												1	1	1																
Hemiptera																																															
Aphididae	A																																1				2	1				4			1		
Coleoptera																																															
Dytiscidae																																															
Hydroporus	L	1		2		1								1												2	2	1																		
Hydroporus	A	1																																													
Gyrinidae																																															
Gyrinus	L				1																						1																			
Ephemeroptera	N		1			20	10		1						20									2		21			1	1	10					5	2	1			10		1			
Baetidae	N						10												1			10	15	2	1	10																			
Procloeon	N				4	4	1	1						2	1					1			10	3	5	10	23	2			7	14	15									1				
Heptageniidae	N																																														
Heptagenia solitaria	N					1																																									
Ephemeridae	N																																														
Ephemera simulans	N																			1		1							1															2	3	4
Caenidae	N																																														
Caenis simulans	N			3	2	2				1				1	1					4	1	1			1	4		3														6	7			1			1
Plecoptera	N																																														
Perlodidae	N					1																																									
Trichoptera	L				2		1		10					1	2							1	2			1																	20				
Hydroptilidae	L																																											4			2
Hydroptila	L														1																															
Leptoceridae	P				1																																									
Mystacides	L																										1																			
Brachycentridae	L																									1																			
Brachycentrus	L																																														
Hemiptera	A	1																															1												
Collembola																																															
Hypogastruridae																																															
Anuridia																					1						10	1								1		5		2						
Odonata																																															
Anisoptera	H					1																																									
Coenagrionidae																																															
Enallagma cyathigerum	H									1																1																					
TOTAL		34	64	53	608	442	326	546	427	1167	273	67	208	780	1090	1059	9	4	3	549	284	652	45	99	40	321	540	210	115	22	91	98	257	266	81	37	18	336	70	54	45	99	0	136	471	314	1142	1777	3841

Table 3. Number of Benthic Invertebrates from Shuswap Lake Foreshore in 2000

Location		12	12	12	12	12	12	12	11	11	11	12	12	12	11	11	11	Total	Sub
Date		July 31	July 31	July 31	July 31	July 31	July 31	July 31	Aug 01	Aug 01	Aug 01	July 31	July 31	July 31	Aug 01	Aug 01	Aug 01		
Sample No.		1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16		
ATS no.		1336	1337	1338	1339	1340	1341	1342	1343	1344	1345	1346	1347	1348	1349	1350	1351		
Substrate		cobble	cobble	cobble	cobble	cobble	sand	sand	sand	sand	cobble	water	water	water	water	water	water		
Recovery Depth (cm)		70	77	68	68	62	69	60	40	31	46								
Species	**Stage**																		
Turbellaria						1													
Nematoda		1	3		1	1		2	1		1								
Tardigrada											1								
Oligochaeta	egg	5		1															
Oligochaeta	juv	3			1			1			3								
Lumbriculidae			1		1	1		1											
Lumbricidae		1																	
Hirudinea																			
Piscicola salmositica	juv						10	5											
Piscicola salmositica		1	1	2		2	2		2	3	4								
Hydracarina		19	10	26	15	28	7	9	4	6	22	1		1					
Oribatidae		1		1		1	3	9		1			1		1				
Forelia		1																	
Ostracoda																			
Candona		1		2	1				1		1	1							
Cycloscypris		2		1	1		5	4			1								
Cypricercus elongata				1			2												
Calanoida																			
Cyclops scutifer				2	3			1											
Eucyclops agilis		2		2	1														
Leptodiaptomus		5		1	5	1	3		7	2	1	5	4	1	7	12	10		
Epischura nevadensis				1			1												
Cyclopoida	cop	7	2	9	6	4	4	5	9	9	27	2	1	1	5				
Harpacticoida																			
Canthocamptidae			1																
Cladocera																			
Daphnidae																			
Daphnia sp.												2				1			
Scapholeberis mucronata				1					3	1	1				4	4			
Macrothricidae																			
Ilyocryptus spinifer			3	2	1														
Chydoridae																			
Acroperus harpae					1														
Alona (costata?)		5	2	1	1	1					1		1			1			
Chydorus sphaericus		1	2	7	2	1	2	1	1		1		1	1	1		1		
Graptoleberis testudinaria			1	1			1								1				
Sididae																			
Sida crystallina		38	26	141	116	96	8	7	34	3	46	8	3	4	8	1			
Amphipoda																			
Hyalella azteca		3		1	1		1			1	3								
Gastropoda	juv/dr	4		6	17	7	9	14	41		10								
Stagnicola		3	2	2	1			1											
Physa										1	1								
Gyraulus		7	1	2	14	12	24	48											
Insecta																			
Chironomidae	L*	2	4	5	4	3			1	2	1								
Chironomidae	P	1					1												
Tanypodinae	P						1												
Paramerina	L			2															
Thienemannimyia	L	10	4		8	2	6	2	1										
Tanytarsini	L*	6	16	4	13	2	3		3	2	18								
Micropsectra	L	6	3		6	2	1		1		7								
Paralauterborniella	L		1		2	1	1				3								
Rheotanytarsus	P										1								
Stempellina	L		1	1	4				7		9								
Stempellina	P			1					2		4								
Tanytarsus	P				1														
Tanytarsus	L				2						2								
Chironomini	L*		1				2			1	5								
Chironomini	P									1									
Demicryptochironomus	L						1		1	2	1			1			1		
Microtendipes	L		1	1					2		11								
Phaenopsectra	L								1		1								
Orthocladiinae	L*	1			4				1	1	1								
Orthocladiinae	P				1	1													
Corynoneura	L	6	3	9		10	2	4	10	4	17			2					
Corynoneura	A	1							1										
Heterotrissocladius	L	1	2		2				2	2	4	1							
Orthocladius	L																		
Psectrocladius	L	1							1		2								
Thienemanniella	L		2		1	1					1								
Diptera																			
Empididae	A	2																	
Coleoptera																			
Dytiscidae																			
Hydroporus	L									1									
Hydroporus	A			1						1	1								
Ephemeroptera	N*			2		2	1		4		7								
Baetidae	N*			1	2	1			4	1	13								
Leptophlebiidae	N*	23	25	29	35	19	12	30	1										
Chorlerpes siblannulata	N	19	11	27	27	29	23	4	4	3	5								
Paraleptophlebia bicornut	N	3	12	4	2		15	3											
Paraleptophlebia tempora	N	2	2	4	1														
Ephemeridae	N																		
Hexagenia limbata	N									1	2								
Caenidae	N																		
Caenis sp.	N						2			1	4								
Trichoptera	L*	2	1	1	7	2	3	1			1		1						
Trichoptera	A																		
Leptoceridae	L*	1			1	1	1	2		1									
Mystacides	L		1						1		1								
Thysanoptera	N						1												
Thysanoptera	A						1												
Collembola																			
Isotoma		14	3	8															
Sminthurus		1																	
Odonata																			
Aeschnidae																			
Aeschna umbrata	N			4	1														
Zygoptera																			
Enallagma cyathigerum	N																		
		212	146	317	313	234	159	154	151	51	246	20	12	11	27	19	12		

Table 2. Number of Benthic Insecta from Shuswap Lake Foreshore in 1999



Table 3. Number of Benthic Invertebrates from Shuswap Lake Foreshore in 2000

Location		12	12	12	12	12	12	12	11	11	11	12	12	12	11	11	11
Date		July 31	July 31	July 31	July 31	July 31	July 31	July 31	Aug 01	Aug 01	Aug 01	July 31	July 31	July 31	Aug 01	Aug 01	Aug 01
Sample No.		1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16
ATS no.		1336	1337	1338	1339	1340	1341	1342	1343	1344	1345	1346	1347	1348	1349	1350	1351
Substrate		cobble	cobble	cobble	cobble	cobble	sand	sand	sand	sand	cobble	water	water	water	water	water	water
Recovery Depth (cm)		70	77	68	68	62	69	60	40	31	46						
Species	**Stage**																
Turbellaria						1											
Nematoda		1	3		1	1		2	1		1						
Tardigrada											1						
Oligochaeta	egg	5		1													
Oligochaeta	juv	3			1			1			3						
Lumbriculidae			1		1	1		1									
Lumbricidae		1															
Hirudinea																	
Piscicola salmositica	juv						10	5									
Piscicola salmositica		1	1	2		2	2		2	3	4						
Hydracarina		19	10	26	15	28	7	9	4	6	22	1		1			
Oribatidae		1		1		1	3	9		1			1		1		
Forelia		1															
Ostracoda																	
Candona		1		2	1				1		1	1					
Cyclocypris		2		1	1		5	4			1						
Cypricercus elongata				1			2										
Calanoida																	
Cyclops scutifer				2	3			1									
Eucyclops agilis		2		2	1												
Leptodiaptomus		5		1	5	1	3		7	2	1	5	4	1	7	12	10
Epischura nevadensis				1			1										
Cyclopoida	cop	7	2	9	6	4	4	5	9	9	27	2	1	1	5		
Harpacticoida																	
Canthocamptidae			1														
Cladocera																	
Daphnidae																	
Daphniasp												2				1	
Scapholeberis mucronata				1					3	1	1				4	4	
Macrothricidae																	
Ilyocryptus spinifer			3	2	1												
Chydoridae																	
Acroperus harpae					1												
Alona (costata?)		5	2	1	1	1					1		1			1	
Chydorus sphaericus		1	2	7	2	1	2	1	1		1		1	1	1		1
Graptoleberis testudinaria			1	1			1								1		
Sididae																	
Sida crystallina		38	26	141	116	98	8	7	34	3	46	8	3	4	8	1	
Amphipoda																	
Hyalella azteca		3		1	1		1			1	3						
Gastropoda	juv/dm	4		6	17	7	9	14	41		10						
Stagnicola		3	2	2	1			1									
Physa										1	1						
Gyraulus		7	1	2	14	12	24	48									
Insecta																	
Chironomidae	L*	2	4	5	4	3			1	2	1						
Chironomidae	P	1					1										
Tanypodinae	P						1										
Paramerina	L			2													
Thienemannimyia	L	10	4		8	2	6	2	1								
Tanytarsini	L*	6	16	4	13	2	3		3	2	18						
Micropsectra	L	6	3		6	2	1		1		7						
Paratanytarsus	L		1		2	1	1				3						
Rheotanytarsus	P										1						
Stempellina	L		1	1	4				2		9						
Stempellina	P			1					2		4						
Tanytarsus	P				1												
Tanytarsus	L				2						2						
Chironomini	L*		1				2			1	5						
Chironomini	P									1							
Demicryptochironomus	L						1		1	2	1			1			1
Microtendipes	L		1	1					2		11						
Phaenopsectra	L								1		1						
Orthocladiinae	L*	1			4				1	1	1						
Orthocladiinae	P				1	1											
Corynoneura	L	6	3	9		10	2	4	10	4	17			2			
Corynoneura	A	1							1								
Heterotrissocladius	L	1	2		2				2	2	4	1					
Orthocladius	L																
Psectrocladius	L	1							1		2						
Thienemanniella	L		2		1	1					1						
Diptera																	
Empididae	A	2															
Coleoptera																	
Dytiscidae																	
Hydroporus	L									1							
Hydroporus	A			1						1	1						
Ephemeroptera	N*			2		2	1		4		7						
Baetidae	N*			1	2	1			4	1	13						
Leptophlebiidae	N*	23	25	29	35	19	12	30	1								
Choroterpes albiannulata	N	19	11	27	27	29	23	4	4	3	5						
Paraleptophlebia bicornut	N	3	12	4	2		15	3									
Paraleptophlebia tempora	N	2	2	4	1												
Ephemeridae	N																
Hexagenia limbata	N									1	2						
Caenidae	N																
Caenis sp	N						2			1	4						
Trichoptera	L*	2	1	1	7	2	3	1			1		1				
Trichoptera	A																
Leptoceridae	L*	1			1	1	1	2		1							
Mystacides	L		1						1		1						
Thysanoptera	N						1										
Thysanoptera	A						1										
Collembola																	
Isotoma		14	3	8													
Sminthurus		1															
Odonata																	
Aeschnidae																	
Aeschna umbrata	N			4	1												
Zygoptera																	
Enallagma cythigerum	N																
		212	148	317	313	234	159	154	151	51	246	20	12	11	27	19	12

Table 4. Number of Non-Insecta Benthic Invertebrates Shuswap Lake 2001

Location Code Name		7	7	7	7	7	7	13	13	13	13	13	13	14	14	14	14	14	14	10	10	10	2	2	2	3	3	3	3	3	3	7	7	7	7	7	7	2		
Date		May 10 day			May 10 night			June 7 day			June 7 night			June 8 night			June 8 day			June 19 day		June 20	July 1 day			July 31 night			July31 day			Aug 27 night			Aug 28 day			Apr 26 day		
Replicate No.		0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36		
Depth (cm)		10	14	27	N/A	N/A	N/A	20	21	19	20	21	19	21	19	19	20	22	18	24	21	17	15	19	15	16	10	12	15	16	14	14	19	20	19	17	17	20		
Volume (mL)		30	30	500	150	150	140	5	5	5	10	5	5	1	1	1	1	1	1	115	30	35	15	10	7	1	1	1	25	10	10	140	160	175	150	250	275	100		
Sample No.		30227	30228	30229	30230	30231	30232	30233	30234	30235	30236	30237	30238	30239	30240	30241	30242	30243	30244	30245	30246	30247	30248	30249	30250	30251	30252	30253	30254	30255	30256	30257	30258	30259	30260	30261	30262	30263	Total	Sub
Species/Group	Stage																																							
COELENTERATA																																								
Hydra												3	1					1		5		20																		
TURBELLARIA														4	11	20				20	6														10					
NEMATODA		1060	1600	970	1260	320	600	29	3	9	8	6	7	3		2	2	5		15		15	12	115	15	3	2	1	3	2	1	100	250	170	160	120	120	120		
TARDIGRADA																							2															10		
OLIGOCHAETA																																								
Enchytraeidae		30		30	30		10	1	9	9	2	2	3				1			135	150	40	1			1	1				3	100	10	50	20	60	30		
Naididae	juv	90	80	120	280	130	50	2	6		1						1			60	50	40	255	470	370	36	11	2	4	4	8	80	160	80	70	80	100	110		
Chaetogaster								1					1	3		1			2	50	25	35	170	100	120	22	13	4	2	6	3	10	20			50	80	80		
Pristina foreli																				30	30		220	225	65		1					40	10		20	10	120	30		
Stylaria lacustris																										2														
Vejdovskyella																										1														
Tubificidae	juv																			7	5											60	90	160	60	120	120	140		
Lumbriculidae				10	1	3	2													1	1					1												10		
Lumbricidae						1																																	1	1623
Hirudinea																																						1		
Dina parva																																1	3	1	3					
Helobdella stagnalis																																3		10	40		40	20	1074	0
Hydracarina	A	9	16	40	150	120	500	11		3	8	7	9	1	6	2	2	2	2	40	5	20	3	1		4			1	1									1	
Frontipoda	A												1					1		5																			1	1001
Torrenticola	A																																40	20	10		80		304	304
Oribatidae	A			20	50	10		1	2		1	2	3	2	3		2	2	2	10	15	5			2		2												37	
OSTRACODA	juv																				5	10																	374	
Candona		30	50	380	10	20		2	1											20	10	10						1					10	10	40	30	80		470	
Cyclocypris				400																	5									1					1	40	20		124	
Limnocythere				100	20	3																																620	1400	
Cypria		40	180	500	3500	2010	7200	31	1		7			3			1					10											60	80	40	10		140	3135	3135
Cypris		1	20	270	960	600	1600	14	2	2										10		10																		
COPEPODA																																							1360	
Cyclopoida		3	2	10	110	30	150	11	4	13	12	14	13	17	30	40	28	25	12	15	10	35	33	110	60	6	7	7	2	2	2	50	40	200	50	70	200	150		
Cyclops capillatus																																30							30	
Harpacticoida																					5	5			10	1	1	1		1			10	20	20	10	40		134	1729
Canthocamptidae				10																																			10	
CLADOCERA																																							0	
Daphnidae																																							10	
Scapholeberis rammneri		430	140					2	3	1								1							10								10			10		200	807	
Leptodoridae																																							0	
Leptodora kindti																						2						1											3	
Chydoridae																																							0	
Acroperus harpae								10	1	8	11	18	36	275	165	245	40	32	33		10	5			15								10	1	20	10	40	10	965	
Alona costata																										4	1		1		1					20			25	
Alonella excisa							10										1	5	2		5		235	270	180	15	2		1	1		50	90	60	310	500	320	10	1567	
Alona guttata																				5	5											40							90	
Chydorus sphaericus					20	10	10	5	4	3	2	4	9	20	3	6	13	32	9			5	2	5	30		1	1	4		1	600	920	380	1310	1300	2060		6369	
Sididae																																							0	
Sida crystallina		1	30					68	23	29	72	146	97	27	13	9	14	14	14	70	70	30		5	5	19	22	20	2			20	20	10			20	670	
Polyphemidae																																							0	
Polyphemus pediculus																		1																				1	13880
AMPHIPODA																																							0	
Hyalella azteca		3	8	10	58	90	100	2		2		2	2							22	8	15				4	1	3	2		1	70	100	70	30	50	100	230	903	903
Gastropoda	juv							1																									10						11	
Lymnaea							1																							1					1			1	3	
Gyraulus					2	1	1							4	2	4	1	2	2	5																			1	
Lymneidae				1																																			1	
Planorbidae		2																												1									43	
Valvata sincera sincera																																							0	
Bivalve																																130	20	40			3	293	
Sphaeriidae	juv	8			1		1																										1	1					2	
Sphaerium																																1			1	1		4	299
Pisidium					1																																		0	
FISH																													1			1							2	
Cottus asper	juv																																						0	
TOTAL		1667	2126	2871	6453	3348	10235	191	59	79	124	204	182	358	233	329	106	125	76	525	438	302	925	1313	870	121	64	40	25	17	17	1289	1974	1323	2244	2452	3604	1912	45771	

Table 5. Number of Benthic Insecta from Shuswap Lake Foreshore in 2001 Method Benthic Pump

Location Code Name		7	7	7	7	7	7	13	13	13	13	13	13	14	14	14	14	14	14	10	10	10	2	2	2	3	3	3	3	3	3	7	7	7	7	7	7	2		
Date		May 10 day			May 10 night			June 7 day			June 7 night			June 8 night			June 8 day			June 19 day		June 20	July 1 day			July 31 night			July 31 day			Aug 27 night			Aug 28 day			Apr 26 day		
Replicate No.		0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36		
Depth (cm)		10	14	27	N/A	N/A	N/A	20	21	19	20	21	19	21	19	19	20	22	18	24	21	17	15	19	15	16	10	12	15	16	14	14	19	20	19	17	17	20		
Volume (mL)		30	30	500	150	150	140	5	5	5	10	5	5	1	1	1	1	1	1	115	30	35	15	10	7	1	1	1	25	10	10	140	160	175	150	250	275	100		
Sample No.		30227	30228	30229	30230	30231	30232	30233	30234	30235	30236	30237	30238	30239	30240	30241	30242	30243	30244	30245	30246	30247	30248	30249	30250	30251	30252	30253	30254	30255	30256	30257	30258	30259	30260	30261	30262	30263	Total	Sub
Species/Group	Stage																																							
Chironomidae	P	3										1														1	1													
Chironomidae	L*															1								10	5															
Tanypodinae	L*																			5																				
Procladius	L																							1																
Thienemannimyia	L							1	2				1							5							1														
Tanytarsini	L*	13	60	100	240	120	160													10				4	5	1	3		1			20	20	20	30	20	40	250		
Cladotanytarsus	L																									1	1	1				8		4	2		40	10		
Cladotanytarsus	P																																				1			
Stempellina	L																										1													
Tanytarsus	L	1	3			2	5			2											1	1	1										1			10		20		
Chironomini	L*	5		60	6	1	2				1	1	3	1							10			4									10			10				
Demicryptochironomus	L					1																	1									1	1	1	1	1				
Microtendipes	L			40	3	1	6																	22	3	1	2			1								20		
Paracladopelma	L																																							
Paracladopelma	P																							3																
Stictochironomus	L	13	9	20	7	5	2																1	1		1						150	70	140	18	60	7	70		
Stictochironomus	P																																	1						
Orthocladiinae	L*	8	30	50	30	20	2				3	5	5	1						5	15	3	2	5	2	3	2						30	20	10	10	20	20		
Orthocladiinae	P																																	2						
Corynoneura	L				1				1			2	2			1		2	1	10	30	1	2		1	6	3	5	10	6	3	50	30	5	50	30	40		
Corynoneura	P																														1			1	1					
Epoicocladius	L																				1																			
Cardiocladius	L																			1													5							
Orthocladius	L				5	2															1		1								2	7	4						
Eukiefferiella	L																			1																				
Paracladius	L																															1								
Psectrocladius	L																						1											1						
Thienemanniella	L																										2	1				20	90	8	30	50	100			
Diamesinae																																								
Potthastia	L					1																	2									1	10							
DIPTERA	L			10																	1																			
Ceratopogonidae																										1														
Bezzia	L												1																						1						
Probezzia	L							9			1		2			1				75	80	15	1		1					1	3				1					
Muscidae																																								
Limnophora	L					1																																		
Tabanidae																																			1					
Tabanus	L																																							
Cecidomyiidae	A	1																																	10					
Empididae	L*																																							
Podonomidae																						1																		
Trichotanypus	L																							1																
Dolichopodidae	A																							2	5															
Rhaphium	L																																							
HYMENOPTERA																						1													20	2	1			
Chalcoidea	A																					1																		
Formicidae	A																																							
Tenthredinidae	A								1							1																								
Aphididae	A									1	1	1												2	2											1	20			
COLEOPTERA																																								
Haliplidae																																								
Haliplus	L						1																																	
Dytiscidae																										1														
Bracebissinus	A				1	4	18					2	2																											
Hydroporus	A		1		2	7	27														2																			
Hydroporus	L																			15	1	5							1											
Oreodytes	L																					1																		
Curculionidae	A																										1													
Staphylinidae	A							1	1											1																	1			
Phalacridae	L																							1																
LEPIDOPTERA	L																																							
TOTAL		44	103	280	295	165	223	11	5	3	6	12	16	2	2	2	2	1	0	128	142	29	16	57	17	16	17	7	12	8	5	256	273	206	165	195	270	391		

Table 4. Number of Non-Insecta Benthic Invertebrates Shuswap Lake 2001

Location Code Name		7	7	7	7	7	7	13	13	13	13	13	13	14	14	14	14	14	14	10	10	10	2	2	2	3	3	3	3	3	3	7	7	7	7	7	7	2
Date		May 10 day			May 10 night			June 7 day			June 7 night			June 8 night			June 8 day			June 19 day		June 20	July 1 day			July 31 night			July 31 day			Aug 27 night			Aug 28 day			Apr 26 day
Replicate No		0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	24	21	17	15	19	15	16	10	12	15	16	14	14	19	20	19	17	17	20
Depth (cm)		10	14	27	N/A	N/A	N/A	20	21	19	20	21	19	21	19	19	20	22	18	24	21	17	15	19	15	16	10	12	15	16	14	140	160	175	150	250	275	100
Volume (mL)		80	80	900	150	150	140	5	5	5	10	5	5	1	1	1	1	1	1	115	80	35	15	10	7	1	1	1	25	10	10	140	160	175	150	250	275	100
Sample No		30227	30228	30229	30230	30231	30232	30233	30234	30235	30236	30237	30238	30239	30240	30241	30242	30243	30244	30245	30246	30247	30248	30249	30250	30251	30252	30253	30254	30255	30256	30257	30258	30259	30260	30261	30262	30263
Species/Group	Stage																																					
COELENTERATA																				5		20																
Hydra												3	1																					10				
TURBELLARIA														4	11	20		1		20	6																	
NEMATODA		1060	1600	970	1260	320	600	29	3	9	8	6	7	3		2	2	5		15			15	12	115	15	3	2	1	3	2	1	100	250	170	160	120	120	120
TARDIGRADA																								2														10
OLIGOCHAETA																																						
Enchytraeidae		30		30	80		10	1	9	9	2	2	3					1		135	150	40	1			3	1					3	100	10	50	20	60	30
Naididae	juv	90	80	120	280	130	50	2	6		1							1		60	50	40	255	470	570	56	11	2	4	4	8	80	160	80	70	80	100	110
Chaetogaster								1					1	5		1			2	90	25	35	170	100	120	22	13	4	2	6	3	10	20			50	80	80
Pristina foreli																				30	30		220	225	65		1					40	10		20	10	120	30
Stylaria lacustris																										2												
Vejdovskiella																										1												
Tubificidae	juv																			7	5											60	80	160	60	120	120	140
Lumbriculidae				10	1	3	2													1	1					1												10
Lumbricidae						1																																1
Hirudinea																																						
Dina parva																																1	3	1	3			
Helobdella stagnalis																																3		10	40		40	20
Hydracarina	A	9	16	40	150	120	900	11			3	8	7	9	1	6	2	2	2	40	1	20	3	1		4			1	1								
Eremigada	A													1					1																		
Terrestricola	A																																					
Oribatidae	A			20	30	10		1	2		1	2	3	2	3		2	2	2	10	15	5		2		2						40	20	10		80	
OSTRACODA	juv																				5	10																
Candona		30	50	350	10	20		2	1											20	30	10					1					10	10	40	30	80	
Cyclocypris				400																	5											10		1	40	20		
Limnocythere				100	20	3																							1									620
Cypria		40	180	500	3500	2010	7200	31	1		7			2			1			10		10										60	80	40	10		140
Cypris		1	20	270	950	680	1600	14	2	2																												
COPEPODA																																						
Cyclopoida		3	2	10	110	30	150	11	4	13	12	14	13	17	30	40	28	25	12	15	10	35	25	110	60	6	7	7	2	2	2	50	40	200	50	70	200	150
Cyclops capillatus																																30						
Harpacticoida																					5	5			10	1	1	1		1			10	20	20	10	40	
Canthocamptidae				10																																		
CLADOCERA																																						
Daphnidae																		1						10								10			10		200
Scapholeberis rammneri		430	140					2	3	1																		1									
Leptodoridae																						2															
Leptodora kindti																																						
Chydoridae																					10	5			15								10	1	20	10	40	10
Acroperus harpae								10	1	8	11	18	36	275	165	245	60	32	33							4	1		1		1				20		
Alona costata																	1	5	2		5		235	270	180	15	2		1	1		50	90	60	310	500	320	10
Alonella excisa							10													5	5											40					
Alona guttata																						5	2	5	50		1	1	4		1	600	920	380	1310	1300	2060	
Chydorus sphaericus					20	10	10	5	4	3	2	4	9	20	3	6	13	32	9																		
Sididae																																20	20	10			20	
Sida crystallina		1	30					65	23	29	72	146	97	27	13	9	14	14	14	70	70	30		5	5	19	22	20	2									
Polyphemidae																		1																				
Polyphemus pediculus																																						
AMPHIPODA																																						
Hyalella azteca		1	8	10	58	90	100	2			2		2	2						22	8	15				4	1	1	2		1	70	100	70	30	50	100	230
Gastropoda	juv							3																									10					
Lymnaea							3																												1			1
Gyraulus					2	1	3							4	2	4	1	2	2	5																		
Lymnaeidae				1																																		
Planorbidae		2																											1									
Valvata sincera sincera																																						
Bivalve																																130	20	40			3	
Sphaeriidae	juv	5			1		1																										1	1				
Sphaerium																																1				1	1
Pisidium					1																																	
FISH																													1			1						
Catostomus	juv																																					
TOTAL		1667	2126	2871	6453	3348	10235	191	59	79	124	204	182	398	233	329	106	125	76	525	438	802	925	1313	870	121	64	40	25	17	17	1289	1974	1323	2244	2452	3604	1912

Table 5. Number of Benthic Insecta from Shuswap Lake Foreshore in 2001 Method Benthic Pump

Location Code Name		7	7	7	7	7	7	13	13	13	13	13	13	14	14	14	14	14	14	10	10	10	2	2	2	3	3	3	3	3	3	7	7	7	7	7	7	2
Date		May 10 day			May 10 night			June 7 day			June 7 night			June 8 night			June 8 day			June 19 day		June 20	July 1 day			July 31 night			July 31 day			Aug 27 night			Aug 28 day			Apr 26 day
Replicate No		0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36
Depth (cm)		10	14	27	N/A	N/A	N/A	20	21	19	20	21	19	21	19	19	20	22	18	24	21	17	15	19	15	16	10	12	15	16	14	14	19	20	19	17	17	20
Volume (mL)		30	30	500	150	150	140	5	5	5	10	5	5	1	1	1	1	1	1	115	30	35	15	10	7	1	1	1	25	10	10	140	160	175	150	250	275	100
Sample No.		30227	30228	30229	30230	30231	30232	30233	30234	30235	30236	30237	30238	30239	30240	30241	30242	30243	30244	30245	30246	30247	30248	30249	30250	30251	30252	30253	30254	30255	30256	30257	30258	30259	30260	30261	30262	30263
Species/Group	Stage																																					
Chironomidae	P	1										1														1	1											
Chironomidae	L*															1								10	5													
Tanypodinae	L*																			5																		
Procladius	L																							1														
Diamesinae/Potthastia	L							1	2				1							5							1											1
Tanytarsini	L*	13	60	100	240	120	160													10				4	5	1	3		1			20	20	20	30	20	40	250
Cladotanytarsini	L																									1	1	1				8		4	2		40	10
Cladotanytarsini	P																																				1	
Stempellina	L																										1											
Tanytarsini	L	1	3			2	5			2																							1					20
Chironomini	L*	5		60	6	1	2				1	1	1	1							10			4										10				
Demicryptochironomus	L					1																		1							1	1	1	1	1			
Microtendipes	L			40	1	1	6																	22	3	1	2			1								20
Parachaetocladius	L																							1														
Parachaetocladius	P																							1		1						150	70	140	18	60	7	70
Stictochironomus	L	13	9	20	7	5	2																											1				
Stictochironomus	P																																					
Orthocladiinae	L*	8	30	50	30	20	2				1	5	5	1						5	15	1	2	5	2	3	2						30	20	10	10	20	20
Orthocladiinae	P																																	2				
Corynoneura	L				1				1			2	2		1		2	1		10	30	1	2		1	6	3	5	10	6	3	50	30	5	50	30	40	
Corynoneura	P																														1			1	1			
Epoicocladius	L																				1																	
Cardiocladius	L																			1													5					
Orthocladius	L				5	2															1			1							2	7	4				
Eukiefferiella	L																			1												1						
Parakiefferiella	L																							1										1				
Psectrocladius	L																										2	1				20	90	8	30	50	100	
Diamesinae	L																																					
Potthastia	L					1																		2								1	10					
DIPTERA	L			10																	1																	
Ceratopogonidae																										1												
Bezzia	L																			75	80	15	1		1					1	1	5			1			
Probezzia	L							9			1			1																									
Muscidae																																						
Limnophora	L					1																																
Tabanidae																																			1			
Tabanus	L																																					
Ceratopogonidae	A	1																																	10			
Empididae	L*																																					
Pelecorhynchidae																						1																
Trichoptera	L																							1														
Dolichopodidae	A																						2	5														
Rhagionidae	L																																					
HYMENOPTERA																						1												20	2	1		
Chalcidoidea	A																					1																
Formicidae	A																																					
Tenthredinidae	A								1							1																			1	20		
Aphididae	A										1	1	1											2	2													
COLEOPTERA																																						
Haliplidae																																						
Haliplus	L						1																															
Dytiscidae																										1												
Bradytarsus	A				1	4	15					2	2																									
Hydroporus	A		1		2	7	27														2																	
Hydroporus	L																			15	1	5							1									
Oreodytes	L																					1																
Curculionidae	A																											1										
Staphylinidae	A							1	1											1																1		
Phalacridae	L																							1														
LEPIDOPTERA	L																																					
TOTAL		44	103	280	295	165	225	11	5	3	6	12	16	2	2	2	2	1	0	178	142	29	16	57	17	16	17	7	12	8	5	256	273	206	165	195	270	391

Table 5. continued

Location Code Name		7	7	7	7	7	7	13	13	13	13	13	13	14	14	14	14	14	14	10	10	10	2	2	2	3	3	3	3	3	3	7	7	7	7	7	7	2		
Date		May 10 day			May 10 night			June 7 day			June 7 night			June 8 night			June 8 day			June 19 day		June 20	July 1 day			July 31 night			July31 day			Aug 27 night			Aug 28 day			Apr 26 day		
Replicate No.		0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36		
Depth (cm)		10	14	27	N/A	N/A	N/A	20	21	19	20	21	19	21	19	19	20	22	18	24	21	17	15	19	15	16	10	12	15	16	14	14	19	20	19	17	17	20		
Volume (mL)		30	30	500	150	150	140	5	5	5	10	5	5	1	1	1	1	1	1	115	30	35	15	10	7	1	1	1	25	10	10	140	160	175	150	250	275	100		
Sample No.		30227	30228	30229	30230	30231	30232	30233	30234	30235	30236	30237	30238	30239	30240	30241	30242	30243	30244	30245	30246	30247	30248	30249	30250	30251	30252	30253	30254	30255	30256	30257	30258	30259	30260	30261	30262	30263	Total	Sub
Species/Group	Stage																																							
EPHEMEROPTERA	N°			1							5	3	1	1				10	5		10	10				2	1							20						
Thysanopetera	A	1												1									1	2																
Baetidae								11		6	6	6	4	6	3	3	3	2	2	80	60	155				14	15	5	1	1	2	12	1	30	1	20	20	130		
Baetis tricaudatus	N																					2												1				1		
Heptageniidae	N°							3	3	2		1	3	8	7	2	4	11	6	50	115	75																		
Cinygmula	N																				10																			
Heptagenein	N																				3																			
Leptophlebiidae																																		21						
Leptophlebia	N				1	1	3	1																														5		
Ephemeridae																																								
Ephemera simulans	N																				2					2						1	10	4	4	1	3		
Caenidae																																								
Caenis simulans	N							5	2	3	3	5	4	3	2	4	4	11	5	1		6																		
Ephemerellidae																																								
Ephemerella inermis	N																	1																						
Tricorythidae																																								
Tricorythodes minutus	N																															3		3			2			
Ameletidae																																								
Ameletus	N	12	12	10	64	50	26	3	1		1		3	1		1		1			2			1																
PLECOPTERA																																								
Suwallia	N																				1																			
TRICHOPTERA	L°																																							
Encyrtidae	A																																	1						
Hydroptilidae	L°																																	20		10	10	20			
Agraylea	P					1																																		
Lepidostomatidae																																								
Lepidostoma	L						1																									4	70	5	3	1	20	10		
Leptoceridae	L°	1																																						
Mystacides	L	2				10	18																			1											3		
Neotrichia	L					2																										11	60	15	2	9	9	3		
Limnephilidae																																								
Onocosmoecus	L					5	3																1		1										1					
HEMIPTERA	N																																							
Corixidae	N°																									4	1	2						1		1	1			
Cortisella	A	3	1		2	8	31															1														1		9		
THYSANOPTERA	N														1								1		1															
COLLEMBOLA																																								
Sminthurus	A			10										2																										
Isotomidae			1		1												1																							
ODONATA																																								
Anisoptera	N°																				5																			
Enallagma	N	1			1																	1																1		
TOTAL		64	118	300	364	242	305	34	11	14	21	27	31	24	15	13	23	32	13	274	345	269	19	60	19	39	34	14	14	9	7	287	414	286	205	239	346	553	5084	

Table 6. Weight (mg) of Benthic Insecta from Shuswap Lake Foreshore in 2001

Location	7	7	7	7	7	7	13	13	13	13	13	13	14	14	14	14	14	14	10	10	10	2	2	2	3	3	3	3	3	3	7	7	7	7	7	7	2
Date	May 10 day			May 10 night			June 7 day			June 7 night			June 8 night			June 8 day			June 19 day		June 20	July 1 day			July 31 night			July31 day			Aug 27 night			Aug 28 day			Apr 26 day
Replicate No.	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36
Depth (cm)	10	14	27	N/A	N/A	N/A	20	21	19	20	21	19	21	19	19	20	22	18	24	21	17	15	19	15	16	10	12	15	16	14	14	19	20	19	17	17	N/A
Volume (mL)	30	30	500	150	150	140	5	5	5	10	5	5	1	1	1	1	1	1	115	30	35	15	10	7	1	1	1	25	10	10	140	160	175	150	250	275	100
Sample No.	30227	30228	30229	30230	30231	30232	30233	30234	30235	30236	30237	30238	30239	30240	30241	30242	30243	30244	30245	30246	30247	30248	30249	30250	30251	30252	30253	30254	30255	30256	30257	30258	30259	30260	30261	30262	30263
Family/Group																																					
Chironomidae	1										0.5												0.1	0.5	0.5	0.01											
Tanypodinae							0.1	0.1				0.1							5.5				0.5			0.2										0.5	
Tanytarsini	1	6.5	10	24	12	0.5			0.2										0.1	0.1	0.5	0.5	0.4	0.1	0.5	0.5	0.2	0.01			4	2	1	1		2	10
Chironomini	15	12	25	12	7	4				0.1	0.1	0.3	0.1							0.1		1	7	1	0.5	0.5			0.2		50	21	7	8	20	40	80
Orthocladiinae	1	3	5	5	3	0.2		0.01		0.3	0.4	0.7	0.1	0.1		0.2	0.1		2	1	0.1	1	0.5	0.3	0.5	0.5	0.3	0.5	0.3	0.4	7	14	2	5	5	8	2
Diamesinae					0.5																	0.1									0.1	1					
Podonomidae																					0.5																
DIPTERA	0.2		5		1					0.5		1		0.5					50	50	4	0.5	8	0.5	0.1				0.5	0.7	0.6			0.5	11.5		
HYMENOPTERA							6	0.01							0.01						2													2	2	2	
COLEOPTERA																																					
Curculionidae																					2																
Dytiscidae		6		22	73	7					7	8							2	8	0.5				4			0.2								
Haliplidae																																					
Phalactidae																																				0.1	
Staphylinidae							0.1	5											0.5							0.2										
LEPIDOPTERA																							0.2														
EPHEMEROPTERA		0.1								0.5	0.3	0.05	0.01			0.1	0.1		0.1						0.1	0.01								0.2			
Ameletidae	2	3	2	14.5	12.5	8	0.1	0.05		0.1		0.3	0.1		0.1		0.1			1			0.2														
Baetidae							0.5		0.3	0.6	0.6	0.4	0.3	0.3	0.2	0.1	0.1	0.1	8	5	20				0.7	0.3	0.5	0.1	0.01	0.02	0.2	0.1	4	0.2	1	1	2
Caenidae							0.5	0.2	0.3	1	1	1	1	0.5	1	1	3	1	0.5	0.5	2																
Ephemerellidae																	7			9																	
Ephemeridae																				56					1						3	44	22	12	1	23	
Heptageniidae							0.1	0.1	0.01		0.1	0.3	0.4	0.7	0.1	0.2	0.5	0.3	5	17	10																
Leptophlebiidae				14	13	44	10																										2			53	
Tricorythidae																															10		9			8	
PLECOPTERA																																					
Chloroperlidae																				6																	
TRICHOPTERA																												0.5									
Hydroptilidae					0.5																													1	1	0.5	
Lepidostomatidae						5																															
Leptoceridae	2				40	21																			0.1						36	120	40	4	29	20	28
Limnephilidae					105	81																													0.5		
HEMIPTERA																																					
Aphididae etc.									0.1	0.1	0.01											0.3	0.2	0.5											0.2	2	
Corixidae	20	5		16	62	232															7				12	0.5	6						6		9	5	69
THYSANOPTERA	0.1												0.1	0.01								0.2	0.2	0.05													
COLLEMBOLA		0.5	0.5	0.2									0.02		0.1																						
ODONATA	4			4															1		1															8	
TOTAL	46.3	36.1	47.5	111.7	329.5	402.7	17.4	5.5	0.9	3.2	10.0	12.2	2.1	2.1	1.5	1.6	10.9	1.4	74.7	153.7	49.6	3.6	17.3	3.0	20.0	2.7	7.0	1.3	1.0	1.1	110.9	202.1	93.0	33.9	80.2	111.6	252.5

Table 5. continued

Location Code Name		7	7	7	7	7	7	13	13	13	13	13	13	14	14	14	14	14	14	19	10	10	2	2	2	3	3	3	3	3	3	7	7	7	7	7	7	2	
Date		May 10 day			May 10 night			June 7 day			June 7 night			June 8 night			June 8 day			June 19 day		June 20	July 1 day			July 31 night			July 31 day			Aug 27 night			Aug 28 day			Apr 26 day	
Replicate No		0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	
Depth (cm)		10	14	27	N/A	N/A	N/A	20	21	19	20	21	19	21	19	19	20	22	18	24	21	17	15	19	15	16	10	12	15	16	14	14	19	20	19	17	17	20	
Volume (mL)		30	30	500	150	150	140	5	5	5	10	5	5	1	1	1	1	1	1	115	30	35	15	10	7	1	1	1	25	10	10	140	160	175	150	250	275	100	
Sample No		30227	30228	30229	30230	30231	30232	30233	30234	30235	30236	30237	30238	30239	30240	30241	30242	30243	30244	30245	30246	30247	30248	30249	30250	30251	30252	30253	30254	30255	30256	30257	30258	30259	30260	30261	30262	30263	
Species/Group	Stage																																						
EPHEMEROPTERA	N*			1							5	3	1	1			10	5		10	10					2	1								20				
Thysanopetera	A	1												1									1	2															
Baetidae								11		6	6	6	4	6	3	3	3	2	2	80	60	155				14	15	5	1	1	2	12	1	30	1	20	20	130	
Baetis tricaudatus	N																					2												1				1	
Heptageniidae	N*							3	3	2		1	3	8	7	2	4	11	6	50	115	75																	
Cinygmula	N																				10																		
Heptagenia	N																				3																		
Leptophlebiidae																																		21					
Leptophlebia	N				1	1	3	1																														5	
Ephemeridae																																							
Ephemera simulans	N																				2					2						1	10	4	4	1	3		
Caenidae																																							
Caenis simulans	N							5	2	3	3	5	4	3	2	4	4	11	5	1		6																	
Ephemerellidae																																							
Ephemerella inermis	N																	1																					
Tricorythidae																																							
Tricorythodes minutus	N																															3		3			2		
Ameletidae																																							
Ameletus	N	12	12	10	64	50	26	3	1		1		3	1		1		1			2			1															
PLECOPTERA																																							
Suwallia	N																				1																		
TRICHOPTERA	L*																												1										
Encyrtidae	A																																	1					
Hydroptilidae	L*																																		10	10	20			
Agraylea	P					1																																	
Lepidostomatidae																																							
Lepidostoma	L						1																																
Leptoceridae	L*	1																														4	70	5	3	1	20	10	
Mystacides	L	2				10	18																			1												3	
Nectopsyche	L					2																										11	60	15	2	9	9	3	
Limnephilidae																																							
Onocosmoecus	L					5	3																																
HEMIPTERA	N																						1		1										1				
Corixidae	N*																									4	1	2						1		1	1			
Corisella	A	3	1		2	8	31															1														1		9	
THYSANOPTERA	N														1								1		1														
COLLEMBOLA																																							
Smanharia	A			10										2																									
Isotomidae			1		1											1																							
ODONATA																																							
Anisoptera	N*																				5																		
Enallagma	N	1			1																	1																1	
TOTAL		64	118	300	364	242	305	34	11	16	21	27	31	24	15	13	23	32	13	274	345	269	19	60	19	39	34	14	14	9	7	287	414	286	205	239	346	553	5084

Table 6. Weight (mg) of Benthic Insecta from Shuswap Lake Foreshore in 2001

Location	7	7	7	7	7	7	13	13	13	13	13	13	14	14	14	14	14	14	10	10	10	2	2	2	3	3	3	3	3	3	7	7	7	7	7	7	2
Date	May 10 day			May 10 night			June 7 day			June 7 night			June 8 night			June 8 day			June 19 day		June 20	July 1 day			July 31 night			July 31 day			Aug 27 night			Aug 28 day			Apr 26 day
Replicate No	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36
Depth (cm)	10	14	27	N/A	N/A	N/A	20	21	19	20	21	19	21	19	19	20	22	18	24	21	17	15	19	15	16	10	12	15	16	14	14	19	20	19	17	17	N/A
Volume (mL)	30	30	500	150	150	140	5	5	5	10	5	5	1	1	1	1	1	1	115	30	35	15	10	7	1	1	1	25	10	10	140	160	175	150	250	275	100
Sample No.	81227	81228	81229	81230	81231	81232	81233	81234	81235	81236	81237	81238	81239	81240	81241	81242	81243	81244	81245	81246	81247	81248	81249	81250	81251	81252	81253	81254	81255	81256	81257	81258	81259	81260	81261	81262	81263
Family/Group																																					
Chironomidae	1										0.5												0.1	0.5	0.5	0.01											
Tanypodinae							0.1	0.1				0.1							5.5				0.5			0.2											0.5
Tanytarsini	1	6.5	10	24	12	0.5			0.2										0.1	0.1	0.5	0.5	0.4	0.1	0.5	0.5	0.2	0.01			4	2	1	1		2	10
Chironomini	15	12	25	12	7	4				0.1	0.1	0.3	0.1							0.1		1	7	1	0.5	0.5			0.2		50	21	7	8	20	40	80
Orthocladiinae	1	3	5	5	3	0.2		0.01		0.3	0.4	0.7	0.1	0.1		0.2	0.1		2	1	0.1	1	0.5	0.3	0.5	0.5	0.3	0.5	0.3	0.4	7	14	2	5	5	8	2
Diamesinae					0.5																	0.1									0.1	1					
Podonomidae																					0.5																
DIPTERA	0.2		5		1					0.5		1		0.5					50	50	4	0.5	8	0.5	0.1				0.5	0.7	0.6			0.5	11.5		
HYMENOPTERA							6	0.01							0.01						2													2	2	2	
COLEOPTERA																																					
Curculionidae																					2																
Dytiscidae		6		22	73	7					7	8							2	8	0.5				4			0.2								
Haliplidae																																					
Phalactidae																																				0.1	
Staphylinidae							0.1	5											0.5							0.2										
LEPIDOPTERA																							0.2														
EPHEMEROPTERA		0.1								0.5	0.3	0.05	0.01			0.1	0.1		0.1						0.1	0.01								0.2			
Ameletidae	2	3	2	14.5	12.5	8	0.1	0.05		0.1		0.3	0.1		0.1		0.1			1			0.2														
Baetidae							0.5		0.3	0.6	0.6	0.4	0.3	0.3	0.2	0.1	0.1	0.1	8	5	20				0.7	0.3	0.5	0.1	0.01	0.02	0.2	0.1	4	0.2	1	1	2
Caenidae							0.5	0.2	0.3	1	1	1	1	0.5	1	1	3	1	0.5	0.5	2																
Ephemerellidae																	7			9																	
Ephemeridae																				56					1						3	44	22	12	1	23	
Heptageniidae							0.1	0.1	0.01		0.1	0.3	0.4	0.7	0.1	0.2	0.5	0.3	5	17	10																
Leptophlebiidae				14	13	44	10																									2					53
Tricorythidae																															10		9			8	
PLECOPTERA																																					
Chloroperlidae																				6																	
TRICHOPTERA																												0.5									
Hydroptilidae					0.5																													1	1	0.5	
Lepidostomatidae						5																															
Leptoceridae	2				40	21																			0.1						36	120	40	4	29	20	28
Limnephilidae					105	81																															
HEMIPTERA																																			0.5		
Apluidae etc									0.1	0.1	0.01											0.3	0.2	0.5										0.2	2	
Corixidae	20	5		16	62	232															7				12	0.5	6						6		9	5	69
THYSANOPTERA	0.1												0.1	0.01								0.2	0.2	0.05													
COLLEMBOLA		0.5	0.5	0.2									0.02		0.1																						
ODONATA	4			4															1		1																8
TOTAL	46.3	36.1	47.5	111.7	329.5	402.7	17.4	5.5	0.9	3.2	10.0	12.2	2.1	2.1	1.5	1.6	10.9	1.4	74.7	153.7	49.6	3.6	17.3	3.0	20.0	2.7	7.0	1.3	1.0	1.1	110.9	202.1	93.0	33.9	80.2	111.6	252.5